



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

C12 N 15/26

(11) Publication number:

**0 173 378**  
A2

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 85201235.0

(22) Date of filing: 25.07.85

(51) Int. Cl.<sup>4</sup>: **C 12 N 15/00**, C 12 N 1/16,  
C 11 D 3/386, C 11 D 3/395,  
C 12 P 21/02, C 07 H 21/04,  
C 12 N 9/02

Best Available Copy

(30) Priority: 27.07.84 EP 84201114  
07.02.85 GB 8503160

(43) Date of publication of application: 05.03.86  
Bulletin 86/10

(84) Designated Contracting States: AT BE CH DE FR GB IT  
LI NL SE

(71) Applicant: UNILEVER NV, Burgemeester  
s'Jacobplein 1 P.O. Box 760, NL-3000 DK Rotterdam (NL)  
(84) Designated Contracting States: BE CH DE FR IT LI NL  
SE AT

(71) Applicant: UNILEVER PLC, Unilever House Blackfriars  
P.O. Box 68, London EC4P 4BQ (GB)

(84) Designated Contracting States: GB

(72) Inventor: Ledebor, Adrianus Marinus, Burg. leFèvre de  
Montignyplein 8, NL-3055 NL Rotterdam (NL)  
Inventor: Maat, Jan, Molenstraat 2, NL-2681 BS Monster  
(NL)  
Inventor: Verrips, Cornelis Theodorus, Hagedoorn 18,  
NL-3142 KB Maassluis (NL)  
Inventor: Visser, Christiaan, Minoeserf 77, NL-2907 VJ  
Capelle a/d IJssel (NL)  
Inventor: Janowicz, Zbigniew Alojzy,  
Adalbert-Stifter-Strasse 49,  
D-4006 Erkrath-Unterefeldhaus (DE)  
Inventor: Hollenberg, Cornelis Petrus, Chopin-Strasse 7,  
D-4000 Düsseldorf (DE)

(74) Representative: van der Toorren, Johannes, Drs. et al,  
UNILEVER N.V. Patent Division Postbus 137, NL-3130 AC  
Vlaardingen (NL)

(54) Use of oxidoreductases in bleaching and/or detergent compositions and their preparation by microorganisms engineered by recombinant DNA technology.

(57) The structural genes and their regulatory DNA sequences of an alcohol oxidase (MOX) and a dihydroxyacetone synthase (DHAS) of *Hansenula polymorpha* have been isolated and the nucleotide sequences determined. The invention relates to the use of the MOX gene, as well as the use of the regulatory DNA sequences of MOX and/or DAS in combination with the MOX gene, optionally after modification thereof, or other oxidase genes, or other genes, to produce engineered microorganisms, in particular yeasts.

Said engineered microorganisms can produce oxidases or other enzymes in yields that allow industrial application on a large scale.

Moreover, said engineered microorganisms can produce oxidases having improved properties with respect to their application in oxidation reactions and/or in bleaching and detergent products.

EP 0 173 378 A2



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

C12 N 15/26

(19)

(11) Publication number:

**0 173 378**  
**A2**

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 85201235.0

(51) Int. Cl.<sup>4</sup>: **C 12 N 15/00**, C 12 N 1/16,  
C 11 D 3/386, C 11 D 3/395,  
C 12 P 21/02, C 07 H 21/04,  
C 12 N 9/02

(22) Date of filing: 25.07.85

No	références, formules, pages à photocopier, etc	No	classement
1		1	C <sub>12</sub> N 15/26
2	512 13 -> EINDE (telle fig.)	2	INF C <sub>12</sub> N 15/78
3	17-18, 23-30, 36-37, Claims Fig 12 C, 14 A, B, C, 16 A -> F	3	INF C <sub>12</sub> N 15/63
4	13-15, Fig 2-4, Claims	4	INF C <sub>12</sub> N 15/82
5	28-30, Claims, Fig 15, 16 A	5	INF C <sub>12</sub> N 15/22
6	28-30, Claims, Fig 15, 16 A	6	INF C <sub>12</sub> N 15/66
7	32, 31-35, 38, Claims	7	INF C <sub>12</sub> N 4/00
8	1-7, 31-38, Claims	8	INF C <sub>12</sub> N 9/04
9	25-27, 38 Claims	9	INF C <sub>12</sub> N 9/06
10	Claims	10	INF C <sub>11</sub> D 3/386

USE OF OXIDOREDUCTASES IN BLEACHING AND/OR DETERGENT  
COMPOSITIONS AND THEIR PREPARATION BY MICROORGANISMS  
ENGINEERED BY RECOMBINANT DNA TECHNOLOGY

The present invention relates to a process for micro-  
biologically preparing oxidoreductases, use of these  
enzymes in bleaching and/or detergent compositions, as  
well as to microorganisms transformed by DNA sequences  
5 coding for an oxidoreductase and optionally for a di-  
hydroxyacetone synthase-enzyme, and H. polymorpha  
alcohol oxidase and/or dihydroxyacetone synthase  
regulation sequences, the microorganisms being suitable  
for use in the process.

10

Oxidoreductases, especially those which use oxygen as  
electron acceptor, are enzymes suitable for use in  
bleaching and/or detergent compositions in which they  
can be used for the in situ formation of bleaching  
15 agents, e.g.  $H_2O_2$ , during the washing or bleaching  
process. See for example

- GB-PS 1 225 713 (Colgate-Palmolive Company), in which  
the use of a mixture of glucose and glucose oxidase  
and other ingredients in a dry powdered detergent  
20 composition has been described,
- DE-PA 2 557 623 (Henkel & Cie GmbH), in which the use  
of a  $C_1$  to  $C_3$  alkanol and alcohol oxidase, or  
galactose and galactose-oxidase, or uric acid and  
uratoxidase, and other ingredients in a dry detergent  
25 composition having bleaching properties has been des-  
cribed, and
- GB-PA 2 101 167 (Unilever PLC) in which the use of a  
 $C_1$  to  $C_4$  alkanol and a  $C_1$  to  $C_4$  alkanol  
oxidase in a liquid bleach and/or detergent com-  
30 position has been described,

wherein the alkanol and the enzyme are incapable of  
substantial interaction until the composition is

diluted with water, and/or has come into contact with sufficient oxygen.

Up to now natural oxidase-enzymes cannot be produced at a cost price that allows industrial application on a large scale, e.g. detergent products. Moreover, the oxidase-enzymes have to act under non-physiological conditions when used in detergent and bleaching products. Further the natural oxidases that have been investigated for use in detergent compositions are accompanied by the natural catalase-enzyme which decomposes almost immediately the peroxide(s) formed, so that no effective bleaching is obtained. Thus a need exists for oxidase-enzymes that are more suitable for use under the conditions of manufacture and use of detergent and bleaching products.

For an economically feasible production of these oxidases it is further required to reach a yield of these enzymes in fermentation processes in the order of that of alcohol oxidase of H. polymorpha, which is up to 20% of the cellular protein (van Dijken et al., 1976).

One way of finding new microorganisms producing enzymes in higher amounts or finding new oxidase-enzymes having improved properties is to check all sorts of microorganisms and try to isolate the relevant oxidases, which are then checked for their abilities to generate peroxides and their stabilities under the conditions of manufacture and use of detergent and bleaching products. One can hope that some day a suitable enzyme will be found, but the chance of success is unpredictable and probably very low.

35

Another way is to apply another trial and error method of crossing the natural microorganisms producing these oxidases by classical genetic techniques, in the hope

that some day one will find a more productive micro-organism or a more suitable enzyme, but again the chance of success is rather low.

5 Clearly, a need exists for a method for preparing  
oxidase-enzymes in higher yield and/or without the con-  
comitant formation of catalase and/or having improved  
properties during storage and/or use in e.g. bleach  
and/or detergent compositions. The problem of trial and  
10 error can be overcome by a process for preparing an  
oxidase-enzyme by culturing a microorganism under  
suitable conditions, and preferably concentrating the  
enzyme and collecting the concentrated enzyme in a  
manner known per se, which process is characterized in  
15 that a microorganism is used that has been obtained by  
recombinant DNA technology and which is capable of pro-  
ducing said oxidase-enzyme.

The microorganisms suitable for use in a process for  
20 preparing an oxidase-enzyme can be obtained by re-  
combinant DNA technology, whereby a microorganism is  
transformed by a DNA sequence coding for an oxidase-  
enzyme (so-called structural gene) together with one or  
more other DNA sequences which regulate the expression  
25 of the structural gene in a particular microorganism or  
group of organisms, either via introduction of an epi-  
somal vector containing said sequences or via a vector  
containing said sequences which is also equipped with  
DNA sequences capable of being integrated into the  
30 chromosome of the microorganism.

The determination of a structural gene coding for the  
enzyme alcohol oxidase (EC 1.1.3.13) originating from  
H. polymorpha together with its regulatory 5'- and 3'-  
35 flanking regions will be described as an example of the  
invention without the scope of the invention being  
limited to this example. The spirit of the invention is

also applicable to the isolation of DNA sequences of other oxidase-enzymes such as glycerol oxidase, glucose oxidase, D-amino acid oxidase etc.; the incorporation of the DNA sequences or modifications thereof into the genome of microorganisms or into episomal vectors used for transforming microorganisms and the culturing of the transformed microorganisms so obtained as such or for producing the desired oxidase-enzymes, as well as the use of these enzymes in bleaching compositions containing them.

Although the microorganisms to be used can be bacteria, e.g. of the genus Bacillus, as well as moulds, the use of yeasts is preferred for technological and economical reasons. In particular a mould or yeast can be selected from the genera Aspergillus, Candida, Geotrichum, Hansenula, Lenzites, Nadsonia, Pichia, Poria, Polyporus, Saccharomyces, Sporobolomyces, Torulopsis, Trichosporon and Zendera, more particularly from the species A. japonicus, A. niger, A. oryzae, C. boidinii, H. polymorpha, Pichia pastoris and Kloeckera sp. 2201. The latter name is sometimes used instead of C. boidinii.

Many  $C_1$ -utilizing yeasts have been isolated during the last decade, and for Hansenula polymorpha and Candida boidinii the methanol metabolism has been studied extensively (for a review see Veenhuis et al., 1983).

The first step in this metabolism is the oxidation of methanol to formaldehyde and  $H_2O_2$  catalysed by MOX. Formaldehyde is oxidized further by the action of formaldehyde dehydrogenase and formate dehydrogenase.  $H_2O_2$  is split into water and oxygen by catalase.

Alternatively, methanol is assimilated into cellular

material. After its conversion into formaldehyde, this product is fixed through the xylulose monophosphate pathway into carbohydrates. Dihydroxyacetone synthase (DHAS) plays a crucial role in this assimilation process.

The appearance of MOX, formate dehydrogenase, formaldehyde dehydrogenase, DHAS and catalase is subject to glucose repression, e.g. on 0.5% glucose. However, synthesis of MOX is derepressed by growth in low concentrations of glucose (0.1%), contrary to the synthesis of DHAS, which is still fully repressed under these conditions (Roggenkamp et al., 1984).

Regulation, i.e. the possibility to switch "on" or "off" of the gene for the polypeptide concerned, is desirable, because it allows for biomass production, when desired, by selecting a suitable substrate, such as, for example melasse, and for production of the polypeptide concerned, when desired, by using methanol or mixtures of methanol and other carbon sources. Methanol is a rather cheap substrate, so the polypeptide production may be carried out in a very economical way.

After derepression of the gene coding for alcohol oxidase (MOX) by growth on methanol, large microbodies, the peroxisomes are formed. While glucose-grown cells contain only a small peroxisome, up to 80% of the internal volume of the cell is replaced by peroxisomes in the derepressed state. The conversion of methanol into formaldehyde and  $H_2O_2$  as well as the degradation of  $H_2O_2$  has been shown to occur in these peroxisomes, while further oxidation or assimilation of formaldehyde most probably occurs in the cytoplasm. This process is a perfect example of compartmentalization of toxic pro-

ducts, of a strong co-ordinate derepression of several cellular processes and of the selective translocation of at least two of the enzymes involved in this process.

5

Most of the enzymes involved in the methanol metabolism have been purified and characterized (Sahm, 1977, Bystrykh et al, 1981). Especially methanol oxidase (EC 1.1.3.13) has been studied in detail. It is an octamer consisting of identical monomers with an  $M_r$  value of about 74 kd and it contains FAD as a prosthetic group. Up to now no cleavable signal sequence for translocation could be detected, as concluded from electrophoresis studies with in vivo and in vitro synthesized products (Roa and Blobel, 1983) or from in vitro synthesis in the presence of microsomal membranes (Roggenkamp et al., 1984).

Under derepressed conditions, up to 20% of the cellular protein consists of MOX.

#### Materials and methods

##### a) Microorganisms and cultivation conditions

Hansenula polymorpha CBS 4732 was obtained from Dr J.P. van Dijken (University of Technology, Delft, The Netherlands). Cells were grown at 37°C in 1 litre Erlenmeyer flasks containing 300 ml minimal medium (Veenhuis et al., 1978), supplemented with 0.5% (v/v) methanol or 0.5% (v/v) ethanol as indicated. Phage lambda L47.1 and the P2 lysogenic E. coli K12 strain Q 364 were obtained from Dr P. van der Elsen (Free University of Amsterdam, The Netherlands) and propagated as described (Loenen and Brammar, 1980).

E. coli K12 strains BHB 2600, BHB 2688 and BHB 2690



(Hohn, 1979) were obtained from Dr M. van Montagu (University of Gent, Belgium), while E. coli K12 strain JM 101.7118 and the M13 derivatives M13 mp 8, 9, 18 and 19 were obtained from Bethesda Research Laboratories Inc. (Gaithersburg, MD, U.S.A.).

b) Enzymes

All enzymes used were obtained from Amersham International PLC, Amersham, U.K., except alpha-helicase which was obtained from Pharm Industrie, Clichy, France. Enzyme incubations were performed according to the instructions of the manufacturer. ATP:RNA adenylyl transferase was purified as described by Edens et al. (1982).

c) Other materials

[<sup>35</sup>S] methionine, [alpha-<sup>35</sup>S] dATP, [alpha-<sup>32</sup>P] dNTP's, [alpha-<sup>32</sup>P] ATP and [gamma-<sup>32</sup>P] ATP were obtained from Amersham International PLC, Amersham, U.K.

Nitrobenzyloxy-methyl (NBM) paper was obtained from Schleicher and Schuell, and converted into the diazo form (DBM) according to the instructions of the manufacturer.

Nitrocellulose filters (type HATF) were obtained from Millipore.

RNA isolation, fractionation and analysis

Hansenula polymorpha cells were grown to mid-exponential phase, either in the presence of methanol or ethanol. The cells were disrupted by forcing them repeatedly through a French Press at 16 000 psi, in a buffer containing 10 mM Tris-HCl pH 8, 5 mM MgCl<sub>2</sub>, 1% NaCl, 6% para-aminosalicylic acid, 1% sodium do-

decylsulphate (SDS) and 5% phenol. The purification of polyadenylated RNA was subsequently performed, as described previously (Edens et al., 1982). One gram cells yielded four mg total RNA and 0.1 mg polyadenylated RNA.

5 Five microgram samples of total RNA or polyadenylated RNA were radioactively labelled at their 3'-ends with ATP:RNA adenylyl transferase and [ $\alpha$ - $^{32}$ P] ATP, and subsequently separated on a 2.5% polyacrylamide gel containing 7 M urea (Edens et al., 1982). For the  
10 preparative isolation of a specific mRNA fraction, 40 micrograms polyadenylated RNA was mixed with four micrograms of labelled polyadenylated RNA and separated on the denaturing polyacrylamide gel. The radioactive 2.4 kb RNA class was eluted from slices of the gel and  
15 freed from impurities by centrifugation through a 5-30% glycerol gradient in 100 mM NaCl, 10 mM Tris-HCl pH 7.5, 1 mM EDTA and 0.1% SDS for 15 h at 24 000 rev./min. in a Beckmann centrifuge using an SW 60 rotor at 20°C. The radioactive fractions were pooled and  
20 precipitated with ethanol. Polyadenylated RNA was translated in vitro in a rabbit reticulocyte lysate according to Pelham and Jackson (1976), using [ $^{35}$ S] methionine as a precursor. The translation products were immuno-precipitated with MOX antiserum as described by Valerio et al. (1983).  
25

#### cDNA synthesis

One third of the RNA fraction, isolated from the polyacrylamide gel, was used to procure a radioactive cDNA with reverse transcriptase (Edens et al., 1982). Using [ $\alpha$ - $^{32}$ P] dATP and [ $\alpha$ - $^{32}$ P] dCTP of a high specific activity (more than 3000 Ci/mM), 20 000 cpm of high molecular weight cDNA was formed during 1 h at  
30 42°C in the presence of human placental ribonuclease inhibitor.  
35

### DNA isolation

Ten g of Hansenula polymorpha cells were washed with 1 M sorbitol and resuspended in 100 ml 1.2 M sorbitol, 10 mM EDTA and 100 mM citric acid pH 5.8, to which 100 microliter beta-mercapto-ethanol was added. Cells were spheroplasted by incubation with 500 mg alpha-helicase for 1 h at 30°C. Spheroplasts were collected by centrifugation at 4000 rev./min. in a Sorvall GSA rotor, resuspended in 40 ml 20 mM Tris-HCl pH 8, 50 mM EDTA and lysed by adding 2.5% SDS. Incompletely lysed cells were pelleted for 30 min. at 20 000 rev./min. in a Sorvall SS34 rotor and DNA was isolated from the viscous supernatant by centrifugation using a CsCl-ethidium bromide density gradient at 35 000 rev./min. for 48 h in a Beckmann centrifuge using a 60 Ti rotor. 2 mg of DNA was isolated with a mean length of 30 kb.

### Preparation of a clone bank in phage lambda L47.1

150 microgram Hansenula polymorpha DNA was partially digested with Sau3AI and sedimented through a 10-40% sucrose gradient in 1 M NaCl, 20 mM Tris-HCl pH 8 and 5 mM EDTA for 22 h at 23 000 rev./min. in an SW 25 rotor. The gradient was fractionated and samples of the fractions were separated on a 0.6% agarose gel in TBE buffer (89 mM Tris, 89 mM Boric acid, 2.5 mM EDTA).

Fractions that contained DNA of 5-20 kb were pooled and the DNA was precipitated with ethanol. Phage lambda L47.1 was grown, and its DNA was isolated as described by Ledebor et al. (1984). The DNA was digested with BamHI and arms were isolated by centrifugation through a potassium acetate gradient as described by Maniatis et al. (1982). Two microgram phage lambda DNA arms and 0.5 µg Sau3AI digested Hansenula polymorpha DNA thus obtained were ligated and packaged in vitro using a

protocol from Hohn (1979). Phages were plated on E. coli strain Q 364 to a plaque density of 20,000 pfu per 14 cm Petri dish. Plaques were blotted onto a nitro-cellulose filter (Benton and Davis, 1977) and the blot  
5 was hybridized with the radioactive cDNA probe isolated as described above. Hybridization conditions were the same as described by Ledebor et al. (1984) and hybridizing plaques were detected by autoradiography.

10 Isolation and partial amino acid sequence analysis of alcohol oxidase (MOX)

Hansenula polymorpha cells grown on methanol were dis-integrated by ultrasonification and the cell debris was  
15 removed by centrifugation. The MOX-containing protein fraction was isolated by  $(\text{NH}_4)_2\text{SO}_4$  precipitation . (40-60% saturation). After dialysis of the precipitate, MOX was separated from catalase and other proteins by  
20 ion-exchange chromatography (DEAE-Sepharose) and gel filtration (Sephacryl S-400). Antibodies against MOX were raised in rabbits by conventional methods using  
complete and incomplete Freund's adjuvants (Difco Lab, Detroit, U.S.A.). Sequence analysis of alcohol oxidase  
25 treated with performic acid was performed on a Beckman sequenator. Identification of the residues was done with HPLC. The amino acid composition was determined on a Chromaspek analyser (Rank Hilger, U.K.), using  
standard procedures and staining by ninhydrine. The  
30 carboxy terminal amino acid was determined as described by Ambler (1972).

Chemical synthesis of deoxyoligonucleotides

Deoxyoligonucleotides were synthesized on a Biosearch  
35 SAM I gene machine, using the phosphite technique (Matteucci and Caruthers, 1981). They were purified on 16% or 20% polyacrylamide gels in TBE.

### Hybridization with deoxyoligonucleotide probes

The deoxyoligonucleotides were radioactively labelled with T<sub>4</sub>-polynucleotide kinase and [ $\gamma$ -<sup>32</sup>P] ATP.

- 5 The DNA of the MOX clones obtained was digested with different restriction enzymes, separated on 1% agarose gel and blotted onto DBM paper. Hybridizations were performed as described by Wallace et al. (1981).

### 10 DNA sequence analysis

- From clone 4 (see Example 1) containing the complete MOX gene, several subclones were made in phage M13mp-8, -9 or M13mp-18, -19 derivatives by standard techniques.
- 15 Small subclones (less than 0.5 kb), cloned in two orientations, were sequenced directly from both sides. From the larger subclones, also cloned in two orientations, sequence data were obtained by an exonuclease Bal31 digestion strategy (see Fig. 1). For each of both
- 20 cloned orientations the RF M13 DNA is digested with a restriction enzyme that preferably cleaves only in the middle of the insert. Subsequently, both orientations of the clones were cut at this unique site, and digested with exonuclease Bal31 at different time intervals.
- 25 Incubation times and conditions were chosen such that about 100-150 nucleotides were eliminated during each time interval. Each fraction was digested subsequently with the restriction enzyme, recognizing the restriction site situated near the position at which the sequence
- 30 reaction is primed in the M13 derivatives. Ends were made blunt end by incubation with T<sub>4</sub>-polymerase and all dNTP's, and the whole mix was ligated under diluted conditions, thereby favouring the formation of internal RF molecules. The whole ligation mix was used to trans-
- 35 form to E. coli strain JM 101-7118. From each time interval several plaques were picked up and sequenced using recently described modifications of the Sanger sequencing protocol (Biggin et al., 1983).

The isolation of auxotrophic mutants

LEU-1 (CBS N° 7171) is an auxotrophic derivative of H. polymorpha strain NCYC 495 lacking  $\beta$ -isopropylmalate  
5 dehydrogenase activity. The isolation of this mutant has been described by Gleeson et al. (1984).

LR9 (CBS N° 7172) is an auxotrophic derivative of H. polymorpha ATCC 34438, lacking orotidine 5'-decarbox-  
10 ylase activity.

For the isolation, all procedures were carried out at 30°C instead of 37°C, which is the optimal temperature for growth of this yeast. Yeast cells were mutagenized  
15 with 3% ethylmethanesulphonate for 2 hr (Fink, 1970). The reaction was stopped with 6% sodium thiosulphate (final concentration) and the solution was incubated for another 10 min. Mutagenized cells were then washed once with H<sub>2</sub>O and incubated for 2 days on YEPD or YNB supple-  
20 mented with uracil for segregation and enrichment of uracil-auxotrophs followed by a 15 hr cultivation on MM without nitrogen source. Finally a nystatin enrichment was employed for 12 hr on MM with a concentration of 10 /ug antibiotic per ml. The treated cells were plated  
25 on YNB plates containing 200 /ug uracil per ml and 0.8 mg 5-fluoroorotic acid (Boeke et al., 1984). Usually 10<sup>6</sup> cells were plated on a single plate. Resistant colonies were picked after 3 days of incu-  
bation, replica plated twice on YNB plates to establish  
30 the auxotrophy. From the auxotrophic mutants ura<sup>-</sup> cells were isolated. Alternatively, 1.5 x 10<sup>6</sup> yeast cells were incubated in one ml of YNB liquid medium supplemented with 200 /ug of uracil and 0.8 mg of 5-fluoroorotic acid. After incubation of 2 days, the  
35 treated cells were plated on YNB containing uracil, replica-plated twice on YNB and analysed as described above.

Such resistant mutants have been shown to be uracil auxotrophs affected at the URA3 or the URA5 locus in S. cerevisiae (F. Lacroute, personal communication). Of about 600 resistant colonies of H. polymorpha tested, 52 exhibited a uracil phenotype. Since URA3 and URA5 mutations in S. cerevisiae lack orotidine 5'-decarboxylase and orotidine 5'-phosphate pyrophosphorylase, respectively (Jones and Fink, 1982), the obtained uracil auxotrophs of H. polymorpha were tested for both enzymatic activities (Lieberman et al., 1955). Mutants affected in either of the two enzymes were found (Table I). They have been designated *odc1* and *oppl* mutants, respectively. The *odc1* mutants exhibit adequate low reversion frequencies (Table II) and thus are suitable for transformation purposes by complementation.

Isolation of autonomous replication sequences (HARS) from H. polymorpha

Chromosomal DNA from H. polymorpha was partially digested either with SalI or BamHI and ligated into the single SalI and BamHI site of the integrative plasmid YIp5, respectively. The ligation mixture was used to transform E. coli 490 to ampicillin resistance. YIp5 is an integrative plasmid containing the URA3 gene as a selective marker (Stinchcomb et al., 1980).

The plasmid pool of H. polymorpha SalI clones was used to transform H. polymorpha mutant LR9. A total of 27 transformants was obtained being also positive in the  $\beta$ -lactamase assay. From all of them, plasmids could be recovered after transformation of E. coli 490 with yeast minilysates. Restriction analysis of the plasmids revealed that most of the inserts show the same pattern. The two different plasmids, pHARS1 and pHARS2, containing inserts of 0.4 and 1.6 kb respectively, were

used for further studies (Fig. 2). Both plasmids transform H. polymorpha mutant LR9 with a frequency of about 500-1,500 transformants per  $\mu$ g of DNA using the transformation procedure of intact cells treated with polyethyleneglycol. Southern analysis of the H. polymorpha transformants after retransformation with pHARS1 and pHARS2 recovered from E. coli plasmid preparations shows the expected plasmid bands and thus excludes integration of the URA3 gene as a cause of the uracil protrophy. Therefore, we conclude that the HARS sequences like ARS1 (Stinchcomb et al., 1982) allow autonomous replication in H. polymorpha. Neither HARS1 nor HARS2 enabled autonomous replication in S. cerevisiae. HARS1 was sequenced completely as shown in Fig. 3.

#### Estimation of plasmid copy number in H. polymorpha transformants

The copy number of plasmids conferring autonomous replication in H. polymorpha either by ARS sequences or by HARS sequences was estimated by Southern blot analysis (Fig. 4). For comparison, plasmid YRP17 in S. cerevisiae (Fig. 4, lanes 6, 7), which has a copy number of 5-10 per cell (Struhl et al., 1979) and the high copy number plasmid pRB58 in S. cerevisiae (Fig. 4, lanes 4, 5) with about 30-50 copies per cell were used. YRP17 is a URA3-containing yeast plasmid, bearing an ARS sequence (Stinchcomb et al., 1982), while pRB58 is a 2  $\mu$ m derivative containing the URA3 gene (Carlson and Botstein, 1982). A Kluyveromyces lactis transformant carrying 2 integrated copies of pBR pBR322 was used as a control (Fig. 4, lanes 2, 3). The intensity of staining in the autoradiogram reveals that the plasmid YRP17 in H. polymorpha has practically the same copy number as in S. cerevisiae, whereas plasmids pHARS-1 and pHARS-2 show a copy number which is in



the range of about 30-40 copies per cell like pBR58 in S. cerevisiae. This proves once more the autonomously replicating character of the HARS sequence.

## 5 Transformation procedures

Several protocols were used.

- 10 a) H. polymorpha strain LEU-1 was transformed using a procedure adapted from Beggs (1978). The strain was grown at 37°C with vigorous aeration in 500 ml YEPD liquid medium up to an OD<sub>600</sub> of 0.5. The cells were harvested, washed with 20 ml distilled water and resuspended in 20 ml 1.2 M sorbitol, 25 mM
- 15 EDTA pH 8.0, 150 mM DTT and incubated at room temperature for 15 minutes. Cells were collected by centrifugation and taken up in 20 ml 1.2 M sorbitol, 0.01 M EDTA, 0.1 M sodium citrate pH 5.8 and 2% v/v beta-glucuronidase solution (Sigma 1500000 units/ml)
- 20 and incubated at 37°C for 105 minutes. After 1 hr, the final concentration of beta-glucuronidase was brought to 4% v/v. For transformation, 3 ml aliquots of the protoplasts were added to 7 ml of ice cold 1.2 M sorbitol, 10 mM Tris-HCl pH 7. Protoplasts
- 25 were harvested by centrifugation at 2000 rpm for 5 minutes and washed three times in ice cold sorbitol buffer. Washed cells were resuspended in 0.2 ml 1.2 M sorbitol, 10 mM CaCl<sub>2</sub>, 10 mM Tris-HCl pH 7 on ice. 2 /ug of YEP13 DNA - an autonomous repli-
- 30 cating S. cerevisiae plasmid consisting of the LEU2 gene of S. cerevisiae and the 2 micron-ori (Broach et al., 1979) - were added to 100 ml of cells and incubated at room temperature. 0.5 ml of a solution of 20% PEG 4000 in 10 mM CaCl<sub>2</sub>, 10 mM Tris-HCl pH
- 35 7.5 was added and the whole mixture was incubated for 2 minutes at room temperature. Cells were collected by brief (5 sec.) centrifugation in an MSG

microfuge set at high speed and resuspended in 0.1 ml YEPD 1.2 M sorbitol pH 7.0, and incubated for 15 minutes at room temperature. The cells were plated directly by surface spreading on plates containing 2% Difco agar, 2% glucose, 0.67% Difco yeast nitrogen base and 20 mg/l of each of L-adenine Hemisulphate, methionine, uracil, histidine, tryptophan, lysine and 1.2 M sorbitol. Leu<sup>+</sup> transformants appear after 5 days incubation at 37°C with a frequency of 50 colonies/ $\mu$ g DNA, while no transformants appear if no DNA is added.

b) Alternatively, H. polymorpha LEU-1 was transformed with YEP13, using a procedure adapted from Das et al. (1984). Exponentially growing cells were grown up to an OD<sub>600</sub> of 0.4, washed in TE buffer (50 mM Tris-HCl pH 8.0, 1 mM EDTA) and resuspended in 20 ml TE buffer. 0.5 ml cells were incubated with 0.5 ml 0.2 M LiCl for 1 hr at 30°C. To 100 ml of these cells 4  $\mu$ g YEP13 in 20 ml TE buffer was added and the sample was incubated for a further 30 minutes at 30°C. An equal volume of 70% v/v PEG 4000 was added and the mixture was incubated for 1 hr at 30°C, followed by 5 min. at 42°C. After addition of 1 ml H<sub>2</sub>O, cells were collected by a brief centrifugation as described under a), washed twice with H<sub>2</sub>O and resuspended in 0.1 ml YEPD 1.2 M sorbitol and incubated for 15 minutes at room temperature. Cells were plated as described. Leu<sup>+</sup> transformants appear with a frequency of 30/ $\mu$ g DNA.

c) The H. polymorpha URA mutant LR9 was transformed with YRP17, a plasmid containing the URA3 gene of S. cerevisiae as a selective marker and an autonomously replicating sequence (ARS) for S. cerevisiae (Stinchomb et al, 1982). Using the protoplast method described by Beggs (1978), 2-5 transformants/ $\mu$ g

DNA were obtained. This number was enlarged, using the  $\text{LiSO}_4$  method of Ito et al. (1983), up to 15-20 transformants per  $\mu\text{g}$  of DNA. However, the best procedure was the procedure described by Klebe et al. (1983), using intact cells treated with PEG 4000. Up to 300 transformants were obtained per  $\mu\text{g}$  DNA. The  $\text{LiSO}_4$  procedure, as well as the Klebe procedure, was performed at  $37^\circ\text{C}$ .

10 Transformation of H. polymorpha based on autonomous replication of the vector was indicated by two characteristics: (1) the instability of the uracil<sup>+</sup> phenotype. After growth of transformants on YEPD for ten generations, more than 99% had lost the ability to grow  
15 on selective medium (Table II). (2) Autonomous replication was further ascertained by transforming E. coli cells with yeast minilysates and retransformation of H. polymorpha. Subsequent Southern analysis showed the presence of the expected plasmid.

20 H. polymorpha LR9 could not be transformed with pRB58, or with pHH85, constructed by insertion of the whole 2 micron circle DNA (Hollenberg, 1982) into the PstI site of the ampicillin gene of plasmid YIP5. YIP5, containing the DNA sequence of HARS1 or HARS2, was transferred to H. polymorpha LR9 using the Klebe protocol  
25 with a frequency of 500-1500 transformants per  $\mu\text{g}$  of DNA. Thus, transformation frequency is 2-5 times higher than described above, using the heterologous ARS 1 in YRP17 of S. cerevisiae. Similarly, the stability of the HARS plasmid in transformants is slightly higher than  
30 the ARS 1 plasmid (Table II).

Transformation of H. polymorpha by integration of the  
35 URA3 gene from S. cerevisiae

The URA3 gene of S. cerevisiae shows no homology to the

ODC gene in H. polymorpha, as revealed by Southern hybridisation of nick-translated YIp5 plasmid DNA to chromosomal DNA of H. polymorpha. Therefore, low-frequency integration of the URA3 gene at random sites of the H. polymorpha genome had to be anticipated. Transformation of mutant LR9 with the integrative vector YIp5 resulted in 30-40 colonies per  $\mu$ g of DNA on YNB plates using the polyethyleneglycol method, whereas no transformants were obtained in the control experiment using YIp5 for transformation of S. cerevisiae mutant YNN27. Analysis of 38 transformants revealed 4 stable integrants after growth on non-selective medium. The integration event was further demonstrated by Southern analysis (Fig. 5).

A second procedure for generating integration of the URA3 gene into chromosomal DNA of H. polymorpha was performed by enrichment of stable Ura<sup>+</sup> transformants from transformants carrying plasmid pHARS1. Transformants were grown in liquid YEPD up to a density of  $10^9$  cells per ml. An aliquot containing  $5 \times 10^6$  cells was used to inoculate 100 ml of fresh medium and was grown up to a cell density of  $10^9$  per ml. The procedure was repeated until about 100 generations had been reached. Since the reversion rate of mutant LR9 is  $2 \times 10^{-9}$  and the frequency of plasmid loss per 10 generations is 97% in pHARS1 transformants, the predominant part of the Ura<sup>+</sup> cells after 100 generations should be integrants. The Ura<sup>+</sup> colonies tested were all shown to maintain a stable Ura<sup>+</sup> phenotype indicating an integration of the URA3 gene. This was further verified by Southern blot analysis. In addition, these data indicate that the integration frequency is  $5 \times 10^{-6}$ .

Example 1CLONING OF THE GENE FOR ALCOHOL OXIDASE (MOX) FROM  
HANSENULA POLYMORPHA5

---

Characterization of polyadenylated RNA

10 Total RNA and polyadenylated RNA, isolated from cells  
grown on methanol, were labelled at their 3'-termini with  
ATP:RNA adenyl transferase, and separated on a de-  
naturing polyacrylamide gel (Fig. 6). Apart from the rRNA  
bands, two classes of RNA appear in the poly-adenylated  
15 RNA lane, respectively 1 kb and 2.3 kb in length. Since  
these RNA classes are not found in polyadenylated RNA of  
ethanol-grown cells (result not shown), they obviously  
are transcripts of genes derepressed by growth on  
methanol. The 2.3 kb class can code for a protein of  
700 to 800 amino acids, depending on the length of the  
20 non-translated sequences. Likewise, the 1 kb class  
codes for a protein of 250-300 amino acids. Enzymes  
that are derepressed by growth on methanol and are 700  
to 800 amino acids long, most likely are MOX (Kato et  
al., 1976; Roa and Blobel, 1983) and DHAS (Bystrykh et  
25 al., 1981). Derepressed enzymes in the 250 to 300 amino  
acid range are probably formaldehyde and formate de-  
hydrogenase (Schütte et al., 1976). The polyadenylated  
RNA was characterized further by in vitro translation  
in a reticulocyte cell free translation system. Two  
30 microliters of the polyadenylated RNA directed protein  
mixture were separated directly on a 10% SDS poly-  
acrylamide gel, while the remaining 18 microliters were  
subjected to immuno-precipitation with antiserum  
against MOX (Fig. 7). Six strong bands dominate in the  
35 total protein mixture, having molecular weights of  
respectively 78kd, 74kd, 58kd, 42kd, 39kd and 36kd.  
Essentially the same molecular weights were found by

Roa and Blobel (1983) in a total cell extract from methanol-grown H. polymorpha cells.

5 The 74kd protein can tentatively be assigned to the monomer of MOX, the 58kd protein to the monomer of catalase and the 39kd and 36kd proteins to the monomers of formaldehyde dehydrogenase and formate dehydrogenase, respectively. The 78kd polypeptide possibly is DHAS, while the 42kd polypeptide remains unidentified.  
10 After immuno-precipitation, both high molecular weight proteins react with the MOX antiserum.

#### Cloning of the gene for MOX

15 Although the 2.3 kb mRNA class induced by growth on methanol obviously codes for at least 2 polypeptides, it seemed a good candidate for screening a Hansenula polymorpha clone bank by hybridization. The 5-20 kb fraction of partially Sau3AI digested H. polymorpha DNA  
20 was cloned in phage lambda L47.1.

Per microgram insert DNA, 300 000 plaques were obtained while the background was less than 1:1000. Two Benton Davis blots, containing about 20 000 plaques each, were  
25 hybridized with 15 000 cpm of the mRNA-derived cDNA probe. After 3 weeks of autoradiography about 40-50 hybridizing plaques could be detected. All plaques were picked up and five were purified further by plating at lower density and by a second hybridization with the  
30 cDNA probe. From four, single hybridizing plaques (1, 3, 4, 5) DNA was isolated. The insert length varied from 8 to 13 kb.

#### Hybridization selection using organic-synthetic DNA probes

35

The sequence of 30 amino acids at the amino terminus of

purified MOX was determined (Fig. 8).

Using the most abundant codon use for the yeast S. cerevisiae, a sequence of 14 bases could be derived  
5 from part of this protein sequence, with only one ambiguity. Both probes, indicated in Fig. 4, were synthesised. In both probes an EcoRI site is present. DBM blots were made from the DNA of the MOX clones digested with the restriction enzymes BamHI, EcoRI/HindIII,  
10 HindIII/SalI and PstI/SalI and separated on 1.5% agarose gels. After hybridization of the blot with a mixture of both radioactively labelled probes, the clones 1, 4 and 5 hybridize, while clone 3 does not, as shown for the HindIII/SalI blot in Fig. 9. However, the  
15 probes did not hybridize with the EcoRI/HindIII digested DNA of these clones (result not shown). Since an EcoRI site is present in the probes, the hybridizing DNA in the clones probably is cut by this enzyme too. Consequently the hybridization overlap has become too  
20 small to allow the formation of stable hybrids.

#### Restriction map and sequence analysis

By comparing restriction enzyme digests and by cross-  
25 hybridization experiments it was concluded that clones 1, 4 and 5 covered identical stretches of DNA.

In order to definitely establish the nature of this stretch of cloned DNA the insert of clone 4 was ana-  
30 lyzed in detail. Hybridization with the amino terminal probe showed that the complete MOX gene (ca. 2 kb) was present, including 2 kb sequences upstream and 3.5 kb downstream (Fig. 10).

35 DNA sequence analysis of the smallest EcoRI fragment revealed the nucleotide sequence corresponding to the amino terminus of MOX as was determined by amino acid sequence analysis.

For sequence analysis, several fragments were subcloned in M13mp8/M13mp9 or M13mpl8/M13mpl9 respectively in two orientations, as indicated in Fig. 10. Clones that were smaller than 0.5 kb were sequenced directly from both  
5 sides. The larger clones were cut at the unique restriction sites situated in the middle of the cloned fragment, to allow generation of exonuclease Bal31 digested subclones as described in materials and methods. Using specific oligonucleotide primers,  
10 sequences around the restriction sites used for subcloning and sequences that did not allow an unequivocal sequence determination were sequenced once more, using the 5.5 kb BamHI/SacI subclone that covers the whole sequence. The complete nucleotide sequence is given in  
15 Fig. 11A and 11B.

The sequence contains an open reading frame of 2046 nucleotides that can code for a protein of 664 amino acids. The last codon of the open reading frame codes  
20 for Phe, which is in agreement with the carboxy terminus of purified MOX. The amino acid composition derived from the DNA sequence encoding this protein, and the amino acid composition of purified MOX are  
virtually identical (Table III). The only important  
25 differences involve the serine and threonine residues, which are notoriously difficult to determine.

The calculated molecular weight of the protein is 74 050 Dalton, which agrees well with the molecular weight of  
30 74 kd of MOX, as determined on polyacrylamide/SDS gels.

#### Codon usage

In Table IV the codon usage for MOX is given. A bias  
35 towards the use of a selective number of codons is evident.



Example 2

CONSTRUCTION OF A PLASMID, pUR 3105, BY WHICH THE GENE  
CODING FOR NEOMYCIN PHOSPHOTRANSFERASE, THAT CONFERS  
5 RESISTANCE AGAINST THE ANTIBIOTIC G 418, IS INTEGRATED  
INTO THE CHROMOSOMAL MOX GENE UNDER REGIE OF THE MOX  
REGULON.

---

10 H. polymorpha cells, transformed with either the plas-  
mids YEP 13, YRP 17, pHARS 1 or pHARS 2, were unstable  
and lost their leu<sup>+</sup> or ura<sup>+</sup> phenotype already after  
10 generations upon growth under non-selective con-  
ditions. In order to obtain stable transformants and to  
15 test the MOX promoter, a plasmid pUR 3105 is construc-  
ted in which the neomycin phosphotransferase gene  
(NEO<sup>R</sup>) is brought under direct control of the MOX  
regulon. The construction is made in such a way that  
the first ATG of the NEO<sup>R</sup> gene is coupled to 1.5 kb  
20 of the MOX regulon. The cloning of such a large regu-  
lon fragment is necessary as shorter fragments, that  
do not contain the -1000 region of the regulon, were  
less efficient.

25 The NEO<sup>R</sup> gene was isolated as a 1.1 kb XmaIII-SalI  
fragment from the transposon Tn5, situated from 35 bp  
downstream of the first ATG up to 240 bp downstream of  
the TGA translational stop codon. To avoid a complex  
ligation mixture, first pUR 3101 is constructed (Fig.  
30 12A), which is a fusion of the far upstream SalI-XmaIII  
(position -1510 to position -1128) fragment of the MOX  
regulon, and the NEO<sup>R</sup> gene, subcloned on M13mp9.  
Another plasmid is constructed, pUR 3102, in which the  
1.5 kb SalI-HgiAI fragment of the MOX gene, that covers  
35 nearly the whole MOX regulon, is ligated to a MOX-  
NEO<sup>R</sup> adapter (Fig. 12B) sequence and cloned in M13-mp9.  
The 1.2 kb XmaIII fragment of this plasmid is cloned in-

to the XmaIII site of pUR 3101, resulting in pUR 3103, which is the exact fusion of the MOX regulon and the NEO<sup>R</sup> gene (Fig. 12C). The orientation is checked by cleavage with HgiAI and SalI. From the lambda-MOX-4  
5 clone, a SalI-SacI fragment is subcloned that reaches from the SalI site, still in the structural MOX gene (position 894), up to the SacI site, far downstream of the structural MOX gene (position 3259) (see Fig. 10). This M13mp19 subclone is called pUR 3104. The plasmid  
10 pUR 3105 is obtained by the direct ligation of the 2.7 kb SalI fragment from pUR 3103 into the SalI site of pUR 3104. The orientation is tested by cleavage with SmaI and SacI.

15 After cleavage of this plasmid with HindIII and SacI and the transformation of this cleaved plasmid to H. polymorpha, G 418-resistant colonies are found that do not lose their resistance upon growth under non-selective conditions for a large number of generations.

Example 3

THE CONSTRUCTION OF pUR 3004, BY WHICH THE GENE CODING  
FOR D-AMINO ACID OXIDASE IS TRANSFERRED TO THE CHROMO-  
5 SOME OF H. polymorpha UNDER REGIE OF THE MOX-REGULON

---

D-amino acid oxidase (AAO) is an example of an oxido-  
reductase for the production of which the methylo-  
10 trophic H. polymorpha is extremely suited. It might be  
expected that the enzyme, being an oxidase like MOX, is  
translocated to the peroxisomes of the yeast that are  
induced during growth on methanol or a mixture of  
methanol and a fermentable sugar as carbon source and  
15 D-amino acids as the sole nitrogen source. Under these  
conditions the cell will be protected from the H<sub>2</sub>O<sub>2</sub>  
produced. Alternatively, AAO can be produced without  
the production of H<sub>2</sub>O<sub>2</sub>, when it is placed under  
regie of the MOX- or DAS-regulon. The AAO production  
20 will be induced by the presence of methanol in the  
medium.

The amino acid sequence of the AAO enzyme has been pub-  
lished (Ronchi et al., 1981) and the complete gene is  
25 synthesised, using the phosphite technique (Matteucci  
and Caruthers, 1981). The gene is constructed in such  
a way that the optimal codon use for H. polymorpha, as  
derived from the sequence of the MOX gene, is used.  
Moreover, several unique restriction sites are intro-  
30 duced without changing the amino acid sequence, to  
facilitate subcloning during the synthesis. The DNA  
sequence is shown in Fig. 13. The gene is synthesised  
in oligonucleotides of about 50 nucleotides in length.  
Oligonucleotides are purified on 16% polyacrylamide  
35 gels. The oligonucleotides that form a subclone are  
added together in ligase buffer (Maniatis et al., 1982)  
and heated to 70°C in a waterbath. The waterbath is

slowly cooled to 16°C and T<sub>4</sub>-ligase is added. After two hours of ligation, the DNA is separated on a 1.5% agarose gel and the fragment, having the expected length, is isolated from the gel. It is subcloned in an M13mp18 vector cleaved at the respective restriction sites situated at the end of the fragment. The gene is subcloned in this way in 4 subclones, respectively SalI-HindIII (position 39-346), HindIII-XmaI (position 346-589), XmaI-KpnI (position 589-721) and KpnI-SalI (position 721-1044). The SalI-HindIII and HindIII-XmaI subclones and the XmaI-KpnI and KpnI-SalI subclones are ligated together as two SalI-XmaI subclones in SalI-XmaI cleaved M13mp18. These two subclones are ligated into a SalI cleaved M13mp8, resulting in pUR 3001 (Figs 13, 14A). The whole sequence is confirmed by the determination of the nucleotide sequence using the modified Sanger dideoxy sequencing technique (Biggin et al., 1983).

The construction of the integrative plasmid, containing the AAO gene is shown in Fig. 14A,B. The nearly complete AAO gene is placed upstream of the MOX termination region, by insertion of the AAO gene-containing SalI fragment of pUR 3001, in the unique SalI site of pUR 3104 (see also Fig. 14A), resulting in pUR 3002. The orientation is checked by cleavage with HindIII. The MOX promoter region is isolated as a 1.4 kb SalI-HgiAI fragment from pUR 3102 (Fig. 14A). This fragment is subsequently placed upstream of the AAO gene in pUR 3002, by ligation to partially SalI-digested pUR 3002 in the presence of the HgiAI-SalI MOX-AAO adapter, shown in Fig. 14A. The orientation of the resulting plasmid pUR 3003 is checked again by cleavage with HindIII. This plasmid is integrated into the MOX gene after cleavage with SacI and transformation to H. poly-morpha cells. Transformants are selected by their ability to grow on D-amino acids as nitrogen source in

the presence of methanol as inducer.

As the selection of cells containing the AAO gene is not simple, another selection marker is introduced. To this end, the S. cerevisiae LEU2 gene is integrated in between the structural AAO gene and the MOX terminator. For this construction, the plasmid pURS 528-03 is used. This plasmid is derived from pURY 528-03 described in European patent application 0096910. The construction is shown in Fig. 14C. The deleted carboxy terminal LEU2 gene sequence of pURY 528-03 was replaced by the complete carboxy terminal LEU2 gene sequence from pYeleu 10 (Ratzkin and Carbon, 1977) and the E. coli lac-lac regulon was eliminated. Subsequently the HpaI-SalI fragment of pURS 528-03 containing the LEU2 gene, is blunt end inserted in the SalI site of pUR 3003, situated in between the AAO structural gene and the MOX terminator. The orientation of the resulting plasmid pUR 3004 can be checked by cleavage with SalI and SacI. pUR 3004 integrates in the chromosomal MOX gene of H. polymorpha after transformation of the SacI-cleaved plasmid to a H. polymorpha leu<sup>-</sup> mutant. Selected leu<sup>+</sup> transformants are integrated in the chromosomal MOX gene, together with the AAO gene.

Example 4

THE CONSTRUCTION OF pUR 3204, pUR 3205, pUR 3210 and  
pUR 3211, BY WHICH THE SMALL PEPTIDE HORMONE, THE HUMAN  
5 GROWTH RELEASING FACTOR, IS EXPRESSED UNDER REGIE OF  
THE MOX-REGULON, EITHER BY INTEGRATION INTO THE CHROMO-  
SOMAL MOX GENE (pUR 3203, pUR 3204), OR BY INTEGRATION  
INTO A HARS1-CONTAINING PLASMID (pUR 3205) OR BY FUSION  
TO THE MOX STRUCTURAL GENE (pUR 3209, pUR 3210 and pUR  
10 3211).

---

Human growth hormone releasing factor (HGRF) is a  
small, 44 amino acids long, peptide, that activates the  
15 secretion of human growth hormone from the pituitary  
glands. HGRF can be used in the diagnosis and treatment  
of pituitary dwarfism in man. Since HGRF has been shown  
to induce growth hormone stimulation in numerous  
species, HGRF might be used in the veterinary field too,  
20 by stimulating growth of animals and increase of milk  
production (Coudé et al., 1984). It is difficult to ob-  
tain HGRF from human sources, but it could very well be  
produced by biotechnological processes, once the gene  
has been cloned and transferred to an appropriate host  
25 organism. Also, as a general example of the production  
of a peptide hormone by H. polymorpha, the gene for  
HGRF is synthesised in the optimal codon use of H.  
polymorpha and brought to expression in several ways.

30 For the construction of pUR 3204 and pUR 3205, the gene  
fragment that codes for the carboxy terminal part of  
the protein is synthesised in DNA oligomers of about 50  
nucleotides in length and subcloned as a HindIII-SalI  
fragment in HindIII-SalI cleaved M13mpl8, resulting in  
35 pUR 3201 (Figs 15, 16A). This HindIII-SalI fragment is  
subsequently inserted upstream of the MOX terminator in  
HindIII-SalI cleaved pUR 3104 (Fig. 16A), resulting in

pUR 3202. The MOX promoter is inserted in front of the HGRF gene, by insertion of the SalI-HgiAI MOX-promoter fragment from pUR 3102 (Fig. 16A) in HindIII cleaved pUR 3202, using a HgiAI-HindIII adapter between the  
5 MOX-promoter and the HGRF gene (Figs 15, 16A). The orientation of the resulting plasmid pUR 3203 is checked by cleavage with SalI and HgiAI. pUR 3203 integrates into the chromosomal MOX gene of H. polymorpha after transformation of the SacI cleaved plasmid. Trans-  
10 formants are selected on immunological activity. pUR 3203 is cleaved with SalI, to insert the SalI-HpaI fragment of pURS 528-03 (Fig. 16B) that contains the LEU2 gene. The orientation of this gene in pUR 3204 is checked by cleavage with HindIII and EcoRI. pUR 3204  
15 integrates into the chromosomal MOX gene of H. polymorpha after transformation of the SacI cleaved plasmid (Fig. 16B) to a leu<sup>-</sup> H. polymorpha mutant. Selection on leu<sup>+</sup> transformants. A plasmid, called pUR 3205, that replicates autonomously in H. polymorpha and contains the HGRF gene, is obtained by insertion of the  
20 EcoRI, partially HindIII cleaved 4 kb long fragment of pUR 3203, containing the HGRF gene inserted in between the MOX-promoter and terminator, into partially HindIII-EcoRI cleaved pHARS1 (Figs 2, 16C). The construction of  
25 pUR 3205 is checked by cleavage with HindIII.

The production of small peptides as HGRF by micro-organisms is often unstable as a result of enzymic degradation (Itakura et al., 1977). Fusion to a protein  
30 like MOX, and subsequent transport to the peroxisomes, could prevent degradation. Therefore, we decided to insert the HGRF gene into the unique KpnI site at position 1775 (amino acid 591, Figs 10, 11) of the MOX structural gene. The HGRF gene is synthesised again in  
35 DNA oligomers of 50 nucleotides in length, but now as two KpnI-HindIII subclones that are cloned as a complete HGRF structural gene in M13mpl9, cleaved with

KpnI (plasmid pUR 3206, Figs 17, 16D). Moreover, the ATG triplet coding for the internal methionine of HGRF at position 27 (Coudé et al., 1984) (position 82 of the DNA sequence) is converted into a TGT triplet coding for cysteine. This does not alter the HGRF activity essentially, and facilitates the cleavage of HGRF from the fusion protein by CNBr cleavage (Itakura et al., 1977). From phage lambda MOX-4 (Fig. 10

SphI (position -491)-KpnI fragment is isolated and inserted into SphI-KpnI cleaved M13mpl9. This results in pUR 3207. pUR 3206 is cleaved with KpnI and the HGRF gene is inserted into the KpnI site of pUR 3207, resulting in pUR 3208. The orientation is checked by direct sequence analysis on the single-stranded DNA of pUR 3208. Subsequently the downstream part of the MOX gene, from the unique KpnI site up to the SacI site, is isolated as a 1.5 kb fragment from phage lambda MOX-4 and inserted into SacI - partially KpnI cleaved pUR 3208. The orientation of the resulting plasmid pUR 3209 is checked by digestion with KpnI. pUR 3209 integrates into the chromosomal MOX gene of H. polymorpha after transformation of the SacI, SphI cleaved plasmid. Selection on immunological activity.

This MOX-HGRF fusion gene is inserted into pHARS1 by isolation of the whole fusion gene from partially HindIII, partially EcoRI cleaved pUR 3209, into EcoRI partially HindIII cleaved pHARS1. This results in pUR 3210, which replicates in H. polymorpha after transformation (Fig. 16E). Alternatively, the LEU2- containing SalI-HpaI fragment of pURS 528-03 is inserted into the blunt-ended KpnI site of the HGRF gene, located at the carboxy terminus of the encoded protein, after partial KpnI cleavage of pUR 3209. The resulting plasmid pUR 3211 integrates into the chromosomal MOX gene of H. polymorpha, after transformation of the SacI, SphI cleaved plasmid (Fig. 16F).



### Discussion

From the length of the open reading frame, from the similarity in the amino acid composition of purified  
5 MOX and the DNA derived protein sequence and from the identical 30 N-terminal amino acids, it is concluded that the complete gene for MOX from the yeast Hansenula polymorpha has been cloned. Its calculated molecular weight agrees well with the molecular weight determined  
10 on SDS polyacrylamide gels. Apart from the coding sequence, more than 1200 bp has been sequenced from both the 5'- and the 3'-non-coding regions, reaching from the SalI site upstream of the coding sequence, up to the SacI site downstream. The gene appears not to be  
15 interrupted with intervening sequences.

The protein is not transcribed in the form of a precursor. Based on the determination of the molecular weight, N-terminal signal sequences could not be  
20 detected in earlier studies of Roa and Blobel (1983) or Roggenkamp et al. (1984) as well. In similar studies, it was suggested that also the rat liver peroxisomal enzymes uricase (Goldman and Blobel, 1978) and catalase (Goldman and Blobel, 1978; Robbi and Lazarow, 1978) do  
25 not contain a cleavable N-terminal signal peptide. However, as discussed by these authors, proteolytic degradation could possibly explain the lack of the detection of such a signal sequence.

30 Our sequence results definitely prove that for translocation of this protein to the peroxisome, a cleavable N-terminal signal sequence is not required. Such a translocation signal may well be situated in the internal sequence of the mature protein, as is the case  
35 for ovalbumine (Lingappa et al., 1979). Inspection of the protein sequence reveals the amino acid sequence Gly X Gly Y Z Gly (amino acids 13-18), which is charac-

teristic for FAD-(flavin adenine dinucleotide)-  
containing enzymes (Ronchi et al., 1981).

5 The isolation of the MOX gene described above gives a  
way how to determine the DNA sequence coding for MOX  
and the amino acid sequence of the MOX enzyme.

10 Similarly, the DNA sequences and amino acid sequences  
belonging to other oxidase-enzymes can be isolated and  
determined. The knowledge of the MOX gene sequence can  
be used to facilitate the isolation of genes coding for  
alcohol oxidases or even other oxidases. By comparing  
the properties and the structure of enzymes one can  
probably establish structure function and activity  
15 relationships. One can also apply methods as site-  
directed mutagenesis, or shortening or lengthening of  
the protein coding sequences, modifying the corres-  
ponding polypeptides, to select oxidase-enzymes with  
improved properties, e.g. with increased alkali  
20 stability, improved production, or oxidase-enzymes  
which need a substrate which is more compatible with  
detergent products.

25 Besides the isolation and characterization of the  
structural gene for MOX from the yeast H. polymorpha,  
also the isolation and characterization of the struc-  
tural gene for DHAS from the yeast H. polymorpha has  
been carried out in a similar way.

30 The DNA sequence of DAS is given in Fig. 18A-18C. A  
restriction map is given in Fig. 19. The amino acid  
composition calculated from the DNA sequence of DAS ap-  
peared to be in agreement with the amino acid com-  
position determined after hydrolysis of purified DHAS.

35 The DHAS enzyme catalyses the synthesis of dihydroxy-  
acetone from formaldehyde and xylulose monophosphate.  
This reaction plays a crucial role in the methanol-

assimilation process (cf. Veenhuis et al., 1983).

As described before, the synthesis of MOX and DHAS is subject to glucose repression. It has now been found  
5 that higher levels of MOX are reached when using glucose/methanol mixtures as substrates instead of 0.5% (v/v) methanol. Under the former conditions up to 30% of the cellular protein consists of MOX, compared with up to 20% under the latter conditions.

10 It was considered that in the regulons of MOX and DAS sequences must exist that play a decisive role in the regulation of repression/derepression by glucose or of the induction by methanol. Some homology therefore  
15 might be expected.

A striking homology of the "TATA-boxes" has been found, both having the sequence CTATAAATA. No other homologies in the near upstream region of the MOX and  
20 DAS regulons have been found. Unexpectedly, a detailed study of both regulons has shown a remarkable homology of the regulons for MOX and DAS in the region about 1000 bp upstream of the translation initiation codon. A practically complete consecutive region of 65 bp in  
25 the regulon of MOX is homologous to a 139 bp region in the DAS regulon, interspersed by several non-homologous regions (see Fig. 20). A similar homology is not found in any other region of both genes, that are over 4 kb in length including their upstream and downstream  
30 sequences. It is suggested that these homologous sequences play a role in the regulation of both genes by glucose and methanol. Transformation studies with vectors containing as regulon the first 500 bp upstream of the ATG of the structural gene of MOX, showed that  
35 this shortened MOX-regulon gave rise to a relatively low expression of the indicator gene beta-lactamase. Indicator genes are genes which provide the yeast with

properties that can be scored easily, e.g. the gene for neomycin phosphotransferase giving resistance to the antibiotic G 418 (cf. Watson et al., 1983) or an auxotrophic marker such as leucin.

5

The fact that the far upstream homologous regions in the MOX and DAS genes have different interruptions and the fact that DAS is repressed at 0.1% glucose and MOX is not, suggest that these homologous regions are of importance to the repression-derepression by glucose and/or the induction of the expression in the presence of methanol. This assumption has been found correct indeed, and the presence or absence of these homologous regions can therefore be important for specific applications. For example, if the -1052 to -987 region of the MOX gene or the -1076 to -937 region of the DAS gene is important for the induction of MOX or DAS by methanol, the presence of these regions is required for the expression of MOX or DAS and/or for the induction of other enzymes by methanol. Another example might be the removal of the regions to avoid repression by glucose, which is needed for the expression of genes coding for proteins other than MOX and DHAS under influence of the MOX and/or DAS regulatory regions with glucose as a carbon source.

25

Thus one aspect of the present invention relates to the isolation and complete characterization of the structural genes coding for MOX and DHAS from the yeast H. polymorpha. It further relates to the isolation and complete characterization of the DNA sequences that regulate the biosynthesis of MOX and DHAS in H. polymorpha, notably the regulons and terminators.

30

Moreover, it relates to combinations of genes coding for alcohol oxidase or other oxidases originating from H. polymorpha strains other than H. polymorpha CBS

35

4732, or Hansenula species other than H. polymorpha, or yeast genera other than Hansenula, or moulds, or higher eukaryotes, with the powerful regulon and terminator of the MOX gene from H. polymorpha CBS 4732. These  
5 combinations may be located on vectors carrying amongst others an autonomously replicating sequence originating from H. polymorpha or related species or minichromosomes containing centromeres, and optionally selection marker(s) and telomeres. These combinations may also be  
10 integrated in the chromosomal DNA of H. polymorpha.

Furthermore it relates to combinations of the powerful regulon or parts of it and terminators of the MOX and/or DAS and - by site-directed mutagenesis or other  
15 methods - changed structural genes coding for alcohol oxidase or another oxidase. These changed structural genes may be located on episomal vectors, in minichromosomes or integrated in the chromosomes of H. polymorpha, H. wingeii, H. anomala, and S. cerevisiae  
20 or in other yeasts.

Besides this, the present invention relates to combinations of the regulon and terminator of the MOX and/or DAS gene of H. polymorpha with structural genes  
25 coding for other proteins than oxidases.

A very important and preferred embodiment of the invention is a process for preparing a polypeptide, such as a protein or an enzyme, by culturing a microorganism  
30 under suitable conditions, optionally concentrating the polypeptide and collecting same in a manner known per se, characterized in that a microorganism is used that has been obtained by recombinant DNA technology and carries a structural gene coding for the polypeptide  
35 concerned, the expression of which is under the control of a regulon, comprising a promoter and at least either the -1052 to -987 region of the MOX gene of Hansenula

polymorpha CBS 4732, or the -1076 to -937 region of the DAS gene of Hansenula polymorpha CBS 4732, or a corresponding region of other methylotrophic moulds or yeasts, or an effective modification of any of these regions.

Surprisingly, it has been observed by the present inventors that the regions concerned, which are shown in Fig. 20 and are referred to herein as the -1000 regions of the MOX and DAS genes, are of crucial importance for the expression of the structural gene concerned. Experiments performed with recombinants containing the MOX regulon from which this region was eliminated showed a low level of expression. Therefore, use of a regulon comprising such -1000 region, or an effective modification thereof, i.e. any modification which does not result in a significant mutilation of the function of said region, makes it possible to realize production of a relatively high amount of the desired polypeptide.

A preferred embodiment of this process according to the invention is characterized in that the structural gene concerned has been provided with one or more DNA sequences coding for amino acid sequences involved in the translocation of the gene product into the peroxisomes or equivalent microbodies of the microbial host. Translocation of the produced polypeptide into the peroxisomes or equivalent microbodies improves their stability, which results in a higher yield. For certain kinds of polypeptides, in particular oxidases, such translocation is imperative for survival of the microbial host, i.e. to protect the host against the toxic effects of the hydrogen peroxide produced when the microbial host cells are growing on the substrate of the oxidase. If the oxidase concerned does not contain addressing signals which are functional in the microbial host used in the production process, one

should provide the structural gene with sequences coding for host specific addressing signals, for example by adding such sequences or by substituting these for the original addressing sequences of the

5 gene. Production of a fused polypeptide, in which the fusion partner carries suitable addressing signals, is another possibility. In case methylotrophic yeasts are used in the production process, it is preferred that the DNA sequences consist of the MOX gene or those parts

10 thereof which are responsible for MOX translocation into the peroxisomes or microbodies.

Finally, this aspect of the present invention is related to the synthesis of MOX originating from H. polymorpha in other yeasts.

15

Some microorganisms with the potential of producing alcohol oxidases are summarized below.

20

Yeasts producing alcohol oxidases  
(Taxonomic division according to Lee and Komagata, 1980)

Group 1 Candida boidinii

25

Group 2a Hansenula philodendra  
Pichia lindnerii  
Torulopsis nemodendra  
" pinus  
30 " sonorensis

Group 2b Candida cariosilignicola

Hansenula glucozyma

" henricii

" minuta

5 " nonfermentans

" polymorpha

" wickerhamii

Pichia pinus

" trehalophila

10

Group 2c Candida succiphila

Torulopsis nitratophila

Group 3 Pichia cellobiosa

15

Group 4 Hansenula capsulata

Pichia pastoris

Torulopsis molischiana

20 Moulds producing alcohol oxidases:

Lenzites trabea

Polyporus versicolor

" obtusus

Poria contigua

25

Among the oxidases other than alcohol oxidases, the most interesting are:

- glycerol oxidase,

- aldehyde oxidase,

30 - amine oxidase,

- aryl-alcohol oxidase,

- amino acid oxidase,

- glucose oxidase,

- galactose oxidase,

35 - sorbose oxidase,

- uric acid oxidase,

- chloroperoxidase, and

- xanthine oxidase.



Combinations of the powerful regulons and terminaters of the MOX and DAS genes from H. polymorpha and structural genes for oxidases may be combined with one or more DNA sequences that enable replication of the

5 structural gene in a particular host organism or group of host organisms, for example autonomously replicating sequences or centromeres (and telomeres) originating from H. polymorpha, to suitable vectors that may be transferred into H. polymorpha and related yeasts or other

10 microorganisms.

H. polymorpha mutants LEU-1 and LR9, mentioned on page 12 of this specification, were deposited at the Centraalbureau voor Schimmelcultures at Delft on 15th

15 July, 1985, under numbers CBS 7171 and CBS 7172, respectively.

The above description is followed by a list of references, claims, Tables, Legends to Figures and Figures.

References

1. GB-PS 1 225 713 (Colgate-Palmolive Company; publ.  
24th March 1971; priority date 19th April 1968).
2. DE-PA 2 557 623 (Henkel & Cie GmbH; publ. 30th June  
1977; priority date 20th December 1975).
3. GB-PA 2 101 167 (Unilever PLC; publ. 12th January  
1983; priority date 7th July 1981).
4. van Dijken, J.P., Otto, R. and Harder, W. (1976),  
Arch.Microb. 111, 137-144.
5. Veenhuis, M., van Dijken, J.P. and Harder, W.  
(1983) in Advances in Microbial Physiology,  
Rose, A.H., Gareth Morris, J. and Tempest, D.W.,  
Eds, Vol. 24, pp 1-82, Academic Press, New York.
6. Roggenkamp, R., Janowicz, Z., Stanikowski, B. and  
Hollenberg, C.P. (1984), Mol.Gen.Genet. 194, 489-  
493.
7. Sahm, H. (1977) in Advances in Microbiol. Engi-  
neering, Ghose, T.K., Fiechter, A. and Blakebrough,  
N., Eds, Vol. 6, pp 77-103, Springer-Verlag,  
Berlin.
8. Bystrykh, L.V., Sokolov, A.P. and Trotsenko, Y.A.  
(1981), FEBS Letters 132, 324-328.
9. Roa, M. and Blobel, G. (1983), Proc.Natl.Acad.Sci.  
USA, 80, 6872-6876.
10. Veenhuis, M., van Dijken, J.P., Pilon, S.A.F. and  
Harder, W. (1978), Arch.Microbiol. 117, 1953-163.

11. Loenen, W.A.M. and Brammar, W.J. (1980), *Gene* 20, 249-259.
- 5 12. Hohn, B. (1979) in *Methods in Enzymology*, Wu, R., Ed., Vol. 68, pp 299-309, Academic Press, New York.
- 10 13. Edens, L., Heslinga, L., Klok, R., Ledeboer, A.M., Maat, J., Toonen, M.Y., Visser, C. and Verrips, C.T. (1982), *Gene* 18, 1-12.
14. Pelham, H.R.B. and Jackson, R.J. (1976), *Eur.J. Biochem.* 67, 247-257.
- 15 15. Valerio, D., Duyvensteijn, M.G.C., Meera Khan, P., Geurts van Kessel, A., de Waard, A. and van der Eb, A.J. (1983), *Gene* 25, 231-240.
- 20 16. Ledeboer, A.M., Verrips, C.T. and Dekker, B.M.M. (1984), *Gene* 30, 23-32.
- 25 17. Maniatis, T., Fritsch, E.F. and Sambrook, J. (1982), *Molecular Cloning*, p 278, Cold Spring Harbor Laboratory Publ., New York.
- 30 18. Benton, W.D. and Davis, R.W. (1977), *Science* 196, 180-182.
19. Ambler, R.P. (1972), *Methods in Enzymology*, Vol. 25, pp 262-272, Academic Press, New York.
20. Matteucci, M.D. and Caruthers, M.H. (1981), *J. Am. Chem. Soc.* 103, 3185-3191.
- 35 21. Wallace, R.B., Johnson, M.J., Hirose, T., Miyake, T., Kawashima, E.H. and Itakura, K. (1981), *Nucl. Acids Res.* 9, 879-894.

22. Biggin, M.D., Gibson, T.J. and Hong, G.F. (1983),  
Proc.Natl.Acad.Sci.USA 80, 3963-3965.
- 5 23. Gleeson, M.A., Waites, M.J. and Sudbery, P.E. (1984),  
in: Microbial growth on C<sub>1</sub> compounds, Eds.  
Crawford, R.L. and Hanson, R.S., Publ. A.S.M.,  
Washington, 228-235.
- 10 24. Fink, G.D. (1970), Methods in Enzymology, Tabor, H.  
and Tabor, C.W., Eds., Vol. 17, pp 59-78, Academic  
Press, New York.
- 15 25. Boeke, J.D., LaCroute, F. and Fink, G.D. (1984), Mol.  
Gen.Genet. 197, 345-346.
26. Jones, E.W. and Fink, G.D. (1982), Cold Spring  
Harbour Monogr.Ser., 11B, 181-299.
- 20 27. Lieberman, I., Kornberg, A. and Simms, E.S. (1955),  
J.Biol.Chem. 215, 403-415.
- 25 28. Stinchcomb, D.T., Thomas, M., Kelly, J., Selker, E.  
and Davis, R.W. (1980), Proc.Natl.Acad.Sci.USA 77,  
4559-4563.
29. Stinchcomb, D.T., Mann, C. and Davis, R.W. (1982),  
J.Mol.Biol. 158. 157-179.
- 30 30. Struhl, K., Stinchcomb, D.T., Scherer, S. and Davis,  
R.W. (1979), Proc.Natl.Acad.Sci.USA 76. 1035-1039.
31. Carlson, M. and Botstein, D. (1982), Cell 28, 145-  
154.
- 35 32. Beggs, J.D. (1978), Nature 275, 104-109.
33. Broach, J.R., Strathern, J.N. and Hicks, J.B. (1979),  
Gene 8, 121-133.

34. Das, S., Kellerman, E. and Hollenberg, C.P. (1984),  
J.Bacteriol. 158, 1165-1167.
- 5 35. Ito, H., Fukuda, Y., Murata, K. and Kimura, A.  
(1983), J.Bacteriol. 153, 163-168.
36. Klebe, R.J., Harriss, J.V., Sharp, Z.D. and Douglas,  
M.G. (1983), Gene 25, 333-341.
- 10 37. Hollenberg, C.P. (1982), Curr.Top.Microbiol.Immunol.  
96, 119-144.
38. Kato, N., Omori, Y., Tani, Y. and Ogata, K. (1976),  
Eur.J.Biochem. 64, 341-350.
- 15 39. Schütte, H., Flossdorf, J., Sahm, H. and Kula, M.R.,  
(1976), Eur.J.Biochem. 62, 151-160.
40. Ronchi, S., Minchiotti, L., Galliano, M., Curti, B.,  
20 Swenson, R.P., Williams, C.H. and Massey, V. (1981),  
J.Biol.Chem. 257, 8824-8830.
41. Ratzkin, B. and Carbon, J. (1977), Proc.Natl.Acad.  
Sci.USA 74, 487-491.
- 25 42. Coudé, F.X., Diaz, J., Morre, M., Roskam, W. and  
Roncucci, R. (1984), Trends in Biotechnology 2, 83-  
88.
- 30 43. Itakura, K., Hirose, T., Crea, R., Riggs, A.D.,  
Heyneker, H.L., Bolivar, F. and Boyer, H.W. (1977),  
Science 198, 1056-1063.
44. Goldman, B.M. and Blobel, G. (1978), Proc.Natl.  
35 Acad.Sci.USA 75, 5066-5070.
45. Robbi, M. and Lazarow, P.B. (1978), Proc.Natl.Acad.  
Sci.USA 75, 4344-4348.

46. Lingappa, V.R., Lingappa, J.R. and Blobel, G.  
(1979), Nature 281, 117-121.
- 5 47. Watson, J.D., Tooze, J. and Kurtz, D.T. (1983),  
Recombinant DNA, A Short Course, page 178, published  
by W.H. Freeman and Company, New York.
48. Lee, J.D. and Komagata, K. (1980), J.Gen.Appl.  
Microbiol. 26, 133-158.

TABLE I

Activities of orotidine 5'-phosphate decarboxylase and  
 5 orotidine 5'-phosphate pyrophosphorylase in H. poly-  
morpha mutants requiring uracil for growth.

Strain/ Genotype	Reversion rate	Activity (%) <sup>a</sup>	
		Orotidine 5'- phosphate decarboxylase	Orotidine 5- phosphate pyrophosphorylase
Wild type	-	100	100
15 LR 9/odc1	$< 2 \times 10^9$	$< 1$	106
MR 7/odc1	$6 \times 10^7$	$< 1$	71
NM 8/odc1	$3 \times 10^8$	$< 1$	105
CLK 55/oppl	n.e. <sup>b</sup>	90	$< 1$
CLK 68/oppl	n.e.	82	$< 1$
20 YNN 27/ura3	n.e.	0	n.e.

Strains were grown in YEPD until late exponential  
 25 phase. Extraction of cells was performed with glass  
 beads using a Braun homogenizer. Protein was estimated  
 by the optical density at 280 nm.

- a) Expressed as the percentage of wild type activity.  
 30 b) Not estimated.

TABLE II

Transformation of uracil-requiring mutants of H. poly-  
morpha

5	Strain	Plasmid	Transformation frequency <sup>a</sup>	Stability <sup>b</sup> (%)	Status of transformed DNA
10	LR 9	YRP17	$2.2 \times 10^2$	<1	Autonomous replication
	LR 9	pHARS1	$1.5 \times 10^3$	2	Autonomous replication
15	LR 9	pHARS2	$4.6 \times 10^2$	1.5	Autonomous replication
	LR 9	YIP5	3 (38) <sup>c</sup>	105	Integration
	LR 9	pRB58	0	-	-
	LR 9	pHH85	0	-	-
20	YNN 27	YIP5	0	-	-

a) Expressed as total number per  $\mu$ g of DNA. Intact  
cells treated with polyethyleneglycol were used for  
transformation as described in Materials and Methods.

b) Expressed as the percentage of remaining uracil  
prototrophs after growth on YEPD for ten generations.

c) Number in parentheses indicates the amount of mini-  
colonies containing free plasmid YIP5.



TABLE III

## Amino acid composition of MOX

5	Amino Acid	DNA sequence	Hydrolysate a)
	PHE	31	32
	LEU	47	49
	ILE	34	34
10	MET	12	11
	VAL	42	43
	SER	43	33 a)
	PRO	43	42
	THR	44	38
15	ALA	47	50
	TYR	27	27
	HIS	19	21
	GLN	13	
	GLU	36	] 51
20	ASN	32	
	ASP	50	] 84
	LYS	35	38
	CYS	13	12
	TRP	10	- b)
25	ARG	36	36
	GLY	50	53

a) Hydrolysis was performed for 24 h.

30 b) Not determined.

TABLE IV

Comparison of preferred codon usage in S. cerevisiae,  
H. polymorpha and E. coli

5		<u>Saccharomyces</u>	<u>Hansenula</u>	<u>E. coli</u>
			MOX	
	ALA	GCU, GCC	GCC	GCC not used, no clear pref.
10	SER	UCU, UCC	UCC, UCG	UCU, UCC
	THR	ACU, ACC	ACC	ACU, ACC
	VAL	GUU, GUC	GUA not used, no clear pref.	GUU, GUA
	ILE	AUU, AUC	AUC, AUU	AUC
15	ASP	GAC	GAC	GAC
	PHE	UUC	UUC	UUC
	TYR	UAC	UAC	UAC
	CYS	UGU	no clear pref.	no clear pref.
	ASN	AAC	AAC	AAC
20	HIS	CAC	CAC	CAC
	GLU	GAA	GAG	GAA
	GLY	GGU	GGC practically not used, no clear pref.	GGU, GGC
	GLN	CAA	CAG	CAG
25	LYS	AAG	AAG	AAA
	PRO	CCA	CCU, CCA	CCG
	LEU	UUG	CUG, CUC	CUG
	ARG	AGA	AGA	CGU

Legends to Figures

5           Fig. 1. The exonuclease Bal31 digestion strategy used  
in sequencing specific MOX subclones. The frag-  
ment X-Y subcloned in M13mp-8 or -9, -18 or -19  
is cut at the unique restriction site Z. The  
DNA molecule is subjected to a time-dependent  
exonuclease Bal31 digestion. The DNA fragment  
situated near the M13 sequencing primer is  
10 removed using restriction enzyme Y; ends are  
made blunt end by incubation with T<sub>4</sub>-DNA  
polymerase and then ligated intramolecularly.  
Phage plaques are picked up after trans-  
formation and the fragment is sequenced from  
15 site Z in the direction of site X. Using the  
M13 derivative with a reversed multiple cloning  
site, the fragment is sequenced from site Z in  
the direction of site X.

20       Fig. 2. Alignment of pHARS plasmids derived by in-  
sertion of HARS fragments into the single Sall  
site of YIp5.

25       Fig. 3. The complete nucleotide sequence of the HARS-1  
fragment.

30       Fig. 4. Estimation of copy number by Southern hybrid-  
ization of H. polymorpha transformants. An  
aliquot of 8 and 16 /ul of each probe was  
electrophoresed. Lane 1, phage lambda DNA di-  
gested with HindIII and EcoRI. Lanes 2,3 trans-  
formant of K. lactis containing two copies of  
integrated plasmid, digested with HindIII (M.  
Reynen, K. Breunig and C.P. Hollenberg, un-  
35 published); lanes 4-7, YNN 27, transformed with  
pRB58 (4-5) and YRP17 (6-7) digested with EcoRI  
respectively; lanes 8,9, LR9 transformed with

YRP17 digested with EcoRI; lanes 10,11, LR9 transformed with pHARS2 digested with HindIII; lanes 12,13, LR9 transformed with pHARS1 digested with EcoRI.

5

Fig. 5. Autoradiogram of Southern blots of DNA from H. polymorpha mutant LR9 transformed by integration of plasmid YIp5. Lane 1, phage lambda DNA, digested both with HindIII and EcoRI; lane 2, pHARS-1, undigested; lanes 3-5 and lanes 6,7 show DNA from 2 different transformants. Lane 3, undigested; lane 4, digested with EcoRI; lane 5, digested with PvuII; lane 6, digested with EcoRI; lane 7, digested with PvuII; lane 8, plasmid YIp5, digested with EcoRI. Nick-translated YIp5 was used as a hybridization probe.

10

15

20

25

30

Fig. 6 Electrophoresis of  $^{32}\text{P}$ -labelled RNA from Hansenula polymorpha, purified once (lane A) or twice (lane B) on oligo(dT)cellulose. Electrophoresis was performed on a denaturing 7 M urea 2.5% polyacrylamide gel. The position of the yeast rRNA's and their respective molecular weights are indicated by 18S and 25S. The 2.3 kb band, that can be seen in lane B, was converted into a cDNA probe which was subsequently used to isolate MOX and DHAS clones from the Hansenula polymorpha clone bank.

35

Fig. 7  $^{35}\text{S}$ -labelled proteins obtained after in vitro translation of methanol derepressed, Hansenula polymorpha mRNA with a rabbit reticulocyte lysate. Either 2 microliters of the total lysate (lane A) or an immuno-precipitate of the remaining 18 microliters using a MOX specific antiserum (lane B) were separated on an 11.5%

SDS-polyacrylamide gel. A mixture of proteins with known molecular weights was used as markers.

- 5 Fig. 8. The N-terminal sequence of purified MOX, as determined on a Beckman sequenator. The two probes that could be derived from the sequence Pro-Asp-Gln-Phe-Asp, using Saccharomyces preferred codons, are indicated.

10

- Fig. 9. Hybridization of a DBM blot of HindIII/SalI cut MOX clones. The DNA was separated on a 1.5% agarose gel (Fig. 9A) and the blot was hybridized to a mixture of both MOX-derived synthetic DNA probes (Fig. 8). Only one band of clones 1, 4 and 5 hybridize (Fig. 9B), indicated by an arrow in Fig. 9A. Lane M: molecular weight markers as indicated. Lane A, B, C and D: clones 1, 3, 4 and 5, respectively. Lane E: lambda L47.1.
- 15
- 20

- Fig. 10. Restriction map for MOX clone 4. Only relevant restriction sites are indicated that have been used for subcloning and sequencing of the MOX gene. The open reading frame, containing the structural MOX sequence, and the M13 subclones made are depicted. Restriction sites used are:
- 25
- B = BamHI, E<sub>I</sub> = EcoRI, E<sub>V</sub> = EcoRV, P = PstI, S<sub>I</sub> = SalI, S<sub>C</sub> = SacI, S<sub>T</sub> = StuI, H = HindIII, S<sub>P</sub> = SphI, K = KpnI, Hg = HgiAI and X = XmaI.
- 30

- Fig. 11A,B. The nucleotide sequence of the MOX structural gene and its 5'- and 3'-flanking sequence.

35

- Fig. 12A,C. The construction of plasmid pUR 3105 by which the neomycin phosphotransferase gene

integrates into the chromosomal MOX gene of  
H. polymorpha.

5 Fig. 12B. Promoter MOX-neomycin phosphotransferase  
adapter fragments.

10 Fig. 13. The DNA sequence of the AAO gene, derived from  
the published amino acid sequence. The gene  
is synthesised in the optimal codon use for  
H. polymorpha in oligonucleotides of about 50  
nucleotides long. Restriction sites, used for  
subcloning are indicated. The HgiAI-SalI frag-  
ment forms the adapter between the structural  
AAO gene and the MOX promoter. The trans-  
lational start codon (met) and stop codon  
15 (\*\*\*) are indicated. The structural sequence  
is numbered from 1 to 1044, while the MOX  
promoter is numbered from -34 to -1.

20 Fig. 14A. The construction of pUR 3003, by which the  
AAO gene integrates into the chromosomal MOX  
gene of H. polymorpha. Selection on activity  
of the AAO gene.

25 Fig. 14B. The construction of pUR 3004, by which the  
AAO gene integrates into the chromosomal MOX  
gene of a H. polymorpha leu<sup>-</sup> derivative.  
Selection on leu<sup>+</sup>.

30 Fig. 14C. The construction of pURS 528-03. Owing to the  
removal of the pCR1 sequence and the double  
lac UV5 promoter, this plasmid is about 2.2  
kb shorter than pURY 528-03.

35 Fig. 15. The DNA sequence of the HGRF gene, derived  
from the published amino acid sequence. The  
gene is synthesised in the optimal codon use

for H. polymorpha in oligonucleotides of about 50 nucleotides long. HgiAI, HindIII and SalI sites are used for subcloning. The HgiAI-HindIII fragment forms the adapter between the structural HGRF gene and the MOX promoter. The translational start codon (met) and stop codon (\*\*\*) are indicated. The structural sequence is numbered from 1 to 140, while the MOX promoter is numbered from -34 to -1.

Fig. 16A. The construction of pUR 3203, by which the gene coding for HGRF integrates into the chromosomal MOX gene of H. polymorpha. Selection on immunological activity of HGRF.

Fig. 16B. The construction of pUR 3204, by which the gene coding for HGRF integrates into the chromosomal MOX gene of a H. polymorpha leu<sup>-</sup> derivative. Selection on leu<sup>+</sup>.

Fig. 16C. The construction of pUR 3205, by which the gene coding for HGRF is inserted into a HARS-1-containing plasmid, which replicates autonomously in H. polymorpha. Selection by transformation of a ura<sup>-</sup> mutant.

Fig. 16D. The construction of pUR 3209, by which the gene coding for HGRF integrates into the chromosomal MOX gene of H. polymorpha, fused to the structural MOX gene. HGRF is cleaved from the fusion protein by CNBr cleavage. Selection on immunological activity of HGRF.

Fig. 16E. The construction of pUR 3210, by which the gene coding for HGRF is inserted into a HARS-1-containing plasmid, fused to the structural MOX gene. Selection as in Fig. 16C.

Fig. 16F. The construction of pUR 3211, by which the gene coding for HGRF integrates into the chromosomal MOX gene of a H. polymorpha leu<sup>-</sup> derivative, fused to the structural MOX gene. Selection on leu<sup>+</sup>.

5

Fig. 17. The DNA sequence of the HGRF gene, derived from the published amino acid sequence. The gene is synthesised as mentioned in Fig. 15, but constructed in such a way that it could be inserted into the unique KpnI site of the structural MOX gene. Therefore it was equipped with KpnI sites on both sides of the gene, and KpnI-HindIII fragments were used for sub-cloning. Synthesis will be as a fusion product to the MOX enzyme. The internal met (ATG) at position 82 is converted into a cys (TGT). Translational start (met) and stop (\*\*\*) codons are indicated.

10

15

20

Fig. 18A,B,C. The nucleotide sequence of the DAS structural gene and its 5'- and 3'-flanking sequence.

25

Fig. 19. Restriction map for the DAS-lambda clone. Only relevant restriction sites are indicated that have been used for subcloning and sequencing of the MOX gene. The open reading frame, containing the structural DAS sequence, and the M13 subclones made, are depicted.

30

Fig. 20. Identical sequences in -1000 region of DAS and MOX genes.



CLAIMS

1. Process for preparing an oxidoreductase by culturing a microorganism under suitable conditions, optionally concentrating the enzyme and collecting the concentrated enzyme in a manner known per se, characterized in that a microorganism is used that has been obtained by recombinant DNA technology, and which is capable of producing the oxidoreductase.
2. Process according to claim 1, characterized in that the microorganism is capable of producing at least one enzyme selected from the group consisting of
  - (1) alcohol oxidases,
  - (2) amine oxidases, including alkylamine oxidase and benzylamine oxidase,
  - (3) amino acid oxidases, including D-alanine oxidase, lysine oxidase,
  - (4) cholesterol oxidase,
  - (5) uric acid oxidase,
  - (6) xanthine oxidase,
  - (7) chloroperoxidase, and
  - (8) aldehyde oxidase.
3. Process according to claim 1 or 2, characterized in that the microorganism is a mould or yeast.
4. Process according to claim 3, characterized in that a mould or yeast is selected from the group consisting of the genera Aspergillus, Candida, Geotrichum, Hansenula, Lenzites, Nadsonia, Pichia, Poria, Polyporus, Saccharomyces, Sporobolomyces, Torulopsis, Trichospora and Zendera.
5. Process according to claim 4, characterized in that the mould or yeast is selected from the species Aspergillus japonicus, Aspergillus niger, Aspergillus oryzae, Candida boidinii, Hansenula anomala, Hansenula

polymorpha, Hansenula wingeii, Kloeckera sp.2201 and Pichia pastoris.

6. Process according to any one of claims 1-5,  
5 characterized in that the microorganism is also capable of producing a dihydroxyacetone synthase enzyme, which promotes the formation of dihydroxyacetone from formaldehyde.
- 10 7. Use of an oxidoreductase prepared by a process as claimed in any one of claims 1-5 in an oxidation process.
- 15 8. Bleaching composition including a fabric-washing detergent composition or hard-surface-cleaning composition having bleach activity, characterized in that it contains an oxidoreductase prepared by a process as claimed in any one of claims 1-5 and a substrate for that oxidoreductase.
- 20 9. Microorganism, obtainable by recombinant DNA technology and being capable of producing an oxidoreductase suitable for use in a process as claimed in claims 1-5.
- 25 10. Microorganism, obtainable by recombinant DNA technology and being capable of producing a dihydroxyacetone synthase-enzyme suitable for use in a process according to claim 6, in addition to being capable of  
30 producing an oxidoreductase.
11. Process for preparing a transformed microorganism as claimed in claim 9, characterized in that a DNA sequence coding for an oxidoreductase together with  
35 one or more other DNA sequences which regulate the expression of the structural gene is introduced into the microorganism via an episomal vector or integration

in the genome, such that the microorganism is capable of producing the oxidoreductase.

12. Process for preparing a transformed micro-  
5 organism as claimed in claim 10, characterized in that  
a DNA coding for a dihydroxyacetone synthase-enzyme  
together with one or more other DNA sequences which  
regulate the expression of the structural gene is  
introduced into the microorganism via an episomal  
10 vector or integration in the genome, such that the  
microorganism is capable of producing the dihydroxy-  
acetone synthase-enzyme (DHAS enzyme).

13. DNA sequence coding for an oxidoreductase,  
15 characterized in that it is obtainable by recombinant  
DNA technology from natural and/or cDNA and/or chemi-  
cally synthesised DNA.

14. DNA sequence according to claim 13, charac-  
20 terized in that it codes for an alcohol oxidase.

15. DNA sequence according to claim 14, charac-  
terized in that it comprises the DNA sequence 1-1992  
(MOX gene) given in Fig. 11A + 11B encoding the poly-  
25 peptide 1-664 (MOX), the amino acid sequence of which  
is given in Fig. 11A + 11B.

16. Combination of DNA sequences comprising a  
structural gene coding for an oxidoreductase and one or  
30 more other DNA sequences which regulate the expression  
of the structural gene in a particular microorganism or  
group of microorganisms.

17. Combination of DNA sequences according to  
35 claim 16, characterized in that it comprises at least  
part of the upstream DNA sequence -1 to about -1500  
given in Fig. 11A and/or at least part of the down-

stream DNA sequence 1993 to about 3260 given in Fig. 11B (regulatory regions of the MOX gene).

18. Combination of DNA sequences according to claim 17, characterized in that it comprises at least the polynucleotide -1052 to -987 of the upstream DNA sequence given in Fig. 11A.

19. Combination of DNA sequences according to claim 17, characterized in that it contains a modified MOX promoter sequence which is obtainable by deletion of at least polynucleotide -1052 to -987 given in Fig. 11A.

20. Combination of DNA sequences according to claim 16, characterized in that it comprises at least part of the upstream DNA sequence -1 to about -2125 given in Fig. 18A + 18B and/or at least part of the downstream DNA sequence 2107 to about 2350 given in Fig. 18C (regulatory regions of the DAS gene).

21. Combination of DNA sequences according to claim 20, characterized in that it comprises at least the polynucleotide -1076 to -937 of the upstream DNA sequence given in Fig. 18A.

22. Combination of DNA sequences according to claim 20, characterized in that it contains a modified DAS promoter sequence which is obtainable by deletion of at least polynucleotide -1076 to -937 given in Fig. 18A.

23. Combination of DNA sequences according to claim 16, characterized in that it comprises a structural gene coding for an oxidoreductase of a higher eukaryote, a mould, or a yeast.

24. Combination of DNA sequences according to claim 23, characterized in that it comprises a structural gene coding for an oxidoreductase of a yeast of the genus Hansenula, preferably of the species H. polymorpha.
25. Combination of DNA sequences according to claim 16, characterized in that the structural gene coding for an oxidoreductase encodes an alcohol oxidase.
26. Combination of DNA sequences according to claim 25, characterized in that the structural gene is the DNA sequence 1-1992 (MOX gene) given in Fig. 11A + 11B encoding the polypeptide 1-664 (MOX), the amino acid sequence of which is given in Fig. 11A + 11B.
27. Combination of DNA sequences according to claim 16, characterized in that it also contains a structural gene coding for DHAS.
28. Combination of DNA sequences according to claim 27, characterized in that it contains a structural gene coding for DHAS having the amino acid sequence as given in Fig. 18B + 18C.
29. Combination of DNA sequences according to any one of claims 16-28, characterized in that the DNA sequences have been modified, while retaining their coding function for an oxidoreductase or for their regulatory functions, by recombinant DNA technology.
30. Combination of DNA sequences according to any one of claims 16-29, characterized in that it contains one or more DNA sequences that enable stable inheritance of said combination in the progeny of any particular host organism.

31. Combination of DNA sequences suitable for the transformation of a microbial host to produce a specific enzyme or other protein which combination of DNA sequences contains a regulon, a structural gene  
5 coding for that specific enzyme or other protein and optionally a terminator, characterized in that a regulon is used selected from the group consisting of at least part of the regulon -1 to about -1500 of the MOX gene given in Fig. 11A or at least part of the  
10 regulon of -1 to about -2125 of the DAS gene given in Fig. 18A and modifications thereof that do not impair the regulon function, and optionally a terminator is used selected from the group consisting of at least part of the terminator 1993 to about 3260 of the MOX  
15 gene given in Fig. 11B or at least part of the terminator of 2110 to about 2350 of the DAS gene given in Fig. 18B and modifications thereof that do not impair the terminator function.
- 20 32. Combination of DNA sequences according to claim 31, characterized in that it is suitable for transformation of a Hansenula yeast, in particular a Hansenula polymorpha.
- 25 33. Combination of DNA sequences according to claim 31, characterized in that it is suitable for transformation of a Saccharomyces yeast, in particular Saccharomyces cerevisiae.
- 30 34. Combination of DNA sequences according to claim 31, characterized in that the structural gene coding for that specific enzyme or other protein contains DNA sequences derived from the structural gene coding for MOX (Fig. 11A + 11B), which modify said specific enzyme  
35 or other protein, without impairing its functions, in such a way that said specific enzyme or other protein is translocated into the peroxisomes or equivalent

microbodies of said microbial host.

35. DNA sequence coding for a dihydroxyacetone synthase-enzyme, characterized in that it is obtainable  
5 by recombinant DNA technology from natural and/or cDNA and/or chemically synthesised DNA.

36. DNA sequence according to claim 35, characterized in that it comprises the DNA sequence 1-2106  
10 (DAS gene) given in Fig. 18B + 18C encoding the polypeptide 1-702 (DHAS), the amino acid sequence which is given in Fig. 18B + 18C.

37. Combination of a DNA sequence coding for a  
15 dihydroxyacetone synthase-enzyme and one or more other DNA sequences which regulate the expression of the structural gene in a particular microorganism or group of microorganisms.

20 38. Combination of DNA sequences according to claim 37, characterized in that it comprises the DNA sequence according to claim 36 (DAS gene) and at least part of the upstream DNA sequence -1 to about -2125 given in Fig. 18A + 18B and/or at least part of the  
25 downstream DNA sequence 2107 to about 2350 given in Fig. 18C (regulatory regions of the DAS gene) and/or at least part of the upstream DNA sequence -1 to about -1500 given in Fig. 11A and/or at least part of the downstream DNA sequence 1993 to about 3260 given in  
30 Fig. 11B (regulatory regions of the MOX gene).

39. Combination of DNA sequences according to claim 38, characterized in that it comprises at least the polynucleotide -1076 to -937 of the upstream DNA  
35 sequence given in Fig. 18A or at least the polynucleotide -1052 to -987 of the upstream DNA sequence given in Fig. 11A, respectively.

40. Process for preparing a polypeptide, such as a protein or an enzyme, by culturing a microorganism under suitable conditions, optionally concentrating the polypeptide and collecting same in a manner known per se, characterized in that a microorganism is used that has been obtained by recombinant DNA technology and carries a structural gene coding for the polypeptide concerned, the expression of which is under the control of a regulon, comprising a promoter and at least either the -1052 to -987 region of the MOX gene of Hansenula polymorpha CBS 4732, or the -1076 to -937 region of the DAS gene of Hansenula polymorpha CBS 4732, or a corresponding region of other methylotrophic moulds or yeasts, or an effective modification of any of these regions.

41. Process according to claim 40, characterized in that the promoter is derived from the yeast Hansenula polymorpha.

42. Process according to claim 40 or 41, characterized in that the microorganism is a mould or yeast.

43. Process according to any of claims 40-42, characterized in that a mould or yeast is selected from the group consisting of the genera Aspergillus, Candida, Geotrichum, Hansenula, Lenzites, Nadsonia, Pichia, Poria, Polyporus, Saccharomyces, Sporobolomyces, Torulopsis, Trichospora and Zendera.

44. Process according to claim 43, characterized in that the mould or yeast is selected from the species Aspergillus japonicus, Aspergillus niger, Aspergillus oryzae, Candida boidinii, Hansenula anomala, Hansenula polymorpha, Hansenula wingeii, Kloeckera sp. 2201 and Pichia pastoris.



45. Process according to claim 44, characterized in that the microorganism is the yeast species Hansenula polymorpha.

5 46. Process according to any of claims 40-45, characterized in that the structural gene concerned has been provided with one or more DNA sequences which translocate the gene product into the peroxisomes or equivalent microbodies of the microbial host.

10


47. Process according to claim 46, characterized in that the DNA sequences concerned consist of the MOX gene or those parts thereof which are responsible for MOX translocation into the peroxisomes or microbodies.

\*\*\*\*\*

**THIS PAGE BLANK (USPTO)**


1/32

Fig. 1



**THIS PAGE BLANK (USPTO)**

Fig. 2



**THIS PAGE BLANK (USPTO)**

Fig. 3 DNA sequence of the autonomous-replicating sequence HARS1 from the methylotrophic yeast *Hansenula polymorpha*. The HARS1 represents a *SalI* fragment comprising 483 nucleotides. The dideoxy-sequencing method was employed.

↑  
(G)TCGACTCCG CGACTCGGCG TTCAC TTTTCG AGCTATTTCAT CAACGCCGGA ATACGTCAGA  
AACAGCCGTG CCCAGGGGAC CAGAAAGCCT ACTGGTGAGT ATGTTCTTTC GTGTGATTTT  
CCGAGGATGA GACGACGATA ACGAGCACAA CTCGGAGTCG GAGGACACGC TTATTGCGTI  
GACGAGCCAC ATCAGCAGGC TGTCAAGACT GAGTATAGGC CACAGAGCTG ATTCTGCTCA  
TACTCAAGAC GTTAGTAAAC TCCGTCTGCC ACAATGCTGA CAGAGTATTA TAATAATAGT  
GAATTACGAA CAATGTAGTC AAAAAAATTT AGTAACAATA TGTCATGATG ACAGATTTGC  
TGAAACCAAGT GAACTCCAAT AAATCCAGCG GCTACCGCAT CCCAAGAGAA ACAGATCAGA  
GGTCTAGGCT TGTTTCAGAG TACTACAAGC TTTCCAGAAC TTAGCAATTC TCAAACGCGG  
TTTG(TCGAC)  
↓  
483


**THIS PAGE BLANK (USPTO)**



4/32

0173378

Fig. 4



1 23 45 67 89 10 11 12 13

**THIS PAGE BLANK (USPTO)**

Fig. 5




1 2 3 4 5 6 7 8

**THIS PAGE BLANK (USPTO)**

6/32

0173378



**THIS PAGE BLANK (USPTO)**

0173378

7/32




Fig. 7

**THIS PAGE BLANK (USPTO)**



8/32

0173378

Fig. 8

NH<sub>2</sub>-Ala-Ile-Pro-Asp-Glu-Phe-Asp-Ile-Ile-Val-Val-Gly-

CCA GAC GAA TTC GA

CCA GAT GAA TTC GA


-Gly-Gly- \* -Thr-Gly-Cys-Cys-Ile-Ala-Gly- \* -Leu-  
-Ala-Asn-Leu-Asp-Asp-Gln-Asn-Leu

**THIS PAGE BLANK (USPTO)**


9/32

0173378

Fig. 9



**THIS PAGE BLANK (USPTO)**



**THIS PAGE BLANK (USPTO)**

11/32

0173378

Fig. 11a

CTCGACGCCC AGAACGATCT CCTCCAGCTG CTCCCGGATC AGCTTGTGGC CCGTAATCG  
 -1301  
 AACCAGGCGC AGCCACGCT CTTCCCGAC CACGCTCCCT CGCGAGCCCA GTTTGTGAAC  
 -1451  
 CAGGTCGTTT AGAACCTCCT GCGCAAAGTC CAGTGTCAUA TGAATGTCTT CCTCCGACCA  
 -1351  
 ATTACGATG TTCTCCAGCA GCCATCTGTC TTTCCAGTAG AACCGTAATC TCTGCTCCTC  
 -1301  
 CTACTGTAC CGCAACAGGT AGTTTCCCTC CCCCCCATA ATGAACACGT TCTTTTCTG  
 -1251  
 CTGGCCTGTC ACCAGCGGGC ACCTCTGGAC GCGCTCGATG AGCCCTTCA GCGGCTCGTA  
 -1201  
 GTACTTGTTC CCTCCCTGTA CCCCCCGCGG GTCACGATAC CCACATAGAG GTCTTTGGCC  
 -1151  
 ATTAGTTTGA TGAGGTGGGG CAGGATGGGC GACTCGGCAT CGAAATTTT GCGTCTGTC  
 -1051  
 TACACTGTGA TGTACCATC GAATGTAATC AGCTGCACCT TCGGATCTGG CATCCTTTTC  
 -1001  
 GAATGGAAGA ACCGCGACAT CTCCAACAGC TGGGCGGTGT TGAGAATCAG CCGCAGCTCC  
 -951  
 TTGAACGAGG GGGCCACAAG CCGGCGTTTG CTGATGGCGC GCGGCTCGTC CTGATGTAC  
 -901  
 AAGGCTTTT CCACAGCCAC TCTCGTGAAG AACCTGCCAA CGCTCGCAAC CAUCTGCACG  
 -851  
 AGCCGAGACA ATTUGGGGGT CCGGCGTTTG CTCATTCAA TGTGTCTGTC GATCAGGACT  
 -751  
 TCGAGCTCGT GCAACATTTT CCGGTAGCGG CGTTTTCCT CAGACTTTAC CATGAGGTCC  
 -701  
 TCCACTCCAG AGATCCGGTT GCTCTTCACC CGGTACAGGA CGAACGCCGT GCGCACCAGG  
 -651  
 CCGTTGATCC ATTCTATGAG GCCATCTCCA CGCTCTTCTT TGAATGCGTA CTCCACTCTG  
 -601  
 TAGCGACTGG ACATCTCGAG ACTGGGCTTG CTGTGCTGGA TGCAGCAATT AATTGTGGC  
 -551  
 GCATGCATCC TTGCACCGGA AGTTTTTAAA ACCCACTCGC TTTAGCCGTC GCGTAAACT  
 -451  
 TCTGAATCTG GCAACTGACG GGCTTCTGCA CCGCAACGG AACTTTTCCC TTCCAGGACC  
 -401  
 CAGCTGGATG GTGTATGTG AGGCTCTGTT TGCTGGGTA GCCTACAACG TGACCTTCC  
 -351  
 TAACCGGAGC GCGCTACCCA CTCTGTCTC TGCTCTGTAC CAGAAAATCA CCAGAGCAGC  
 -301  
 ACAGGCGCGA TCTGGCAACT CGTGGGCTGT CCGACAGCGT GTTGTCCAC AGTGCAAAATG  
 -251  
 CCGGTCGAAC GGCAGAAAG TAAATTCTTA TGCTAGCGTG CAGCGACTCC GACATCCCCA  
 -151  
 GTTTTCCCC TACTTCATCA CAGATGGGCT CAGCGCTGCC GCTAAGTCTA CCAACCGTC  
 -101  
 CCCACACGGT CCATCTATAA ATACTGCTCG CACTCCAGGG TCCTCACATC AATCTAAACT  
 -51

1	5	10	15
ACAAAAACAAA	MET ALA ILE PRO ASP GLU PHE ASP ILE ILE VAL VAL GLY CLY CLY SER THR	ATC GCC ATT CTT GAG GAA TTC CAT ATC ATT GTT GTT GGT GGA CGT TCC ACC	
-11			
20	25	30	35
GLY CYS CYS ILE ALA CLY ARG LEU ALA ASN LEU ASP ASP CLN ASN LEU THR VAL ALA LEU	TGC TGC ATT GCG GGC AGA CTG GCA AAC CTC GAC CAA AAC CTC ACA GTT GCC CTG		
40	45	50	55
ILE GLU GLY CLY GLU ASN ASN ILE ASN ASN PRO TRP VAL TYR LEU PRO GLY VAL TYR PRO	ATC CAG GCT CTT CAC AAG AAG ATC AAG AAC CCT TGG GTC TAC CTT CCC GGA CTG TAT CCT		
60	65	70	75
ARG ASN MET ARG LEU ASP SER LYS THR ALA THR PHE TYR SER SER ARG PRO SER LYS ALA	AGA AAC ATG AGA CTC CAC TCC AAC AGC GCC ACC TTC TAC TCC TCC AGA CCA TCC AAG CCT		
80	85	90	95
LEU ASN CLY ARG ARG ALA ILE VAL PRO CYS ALA ASN ILE LEU CLY GLY GLY SER SER ILE	CTG AAC GGC AGA ACA CCG ATC GTT CTT TGC GCC AAC ATC CTT GGA GGC GGC TCC TCC ATC		
100	105	110	115
ASN PHE LEU MET TYR TUR ARG ALA SER ALA SER ASP TYR ASP ASP TRP GLU SER GLU GLY	AAC TTT CTG ATG TAC ACC AGA GCC TCT GGT TCC GAC TAC GAC GAC TCG GAG TCC GAG GGA		
120	125	130	135
TRP SER THR ASP GLU LEU LEU PRO LEU ILE LYS LYS ILE CLU THR TYR CLN ARG PRO CYS	TGG AGC ACC CAC GAG TTG CTA CCT CTG ATC AAA AAA ATC GAA ACT TAC CAG CGT CCT TCC		
140	145	150	155
ASN ASN ARG ASP LEU MET CLY PHE ASP GLY PRO ILE LYS VAL SER PHE GLY ASN TYR THR	AAC AAC AGA CAT CTG CAC GCC TTT GAC GGC CCA ATC AAG GTT TCC TTT GGA AAC TAC ACC		
160	165	170	175
TYR PRO THR CYS CLN ASP PHE LEU ARG ALA ALA GLU SER GLN CLY ILE PRO VAL VAL ASP	TAT CCT ACC TCC CAG CAC TTG CTG ACA GCA GCA CAG TCG GAG GGA ATT CCT GTT GTG GAC		
180	185	190	195
ASP LEU GLU ASP PHE LYS THR SER MET GLY ALA GLU MET TRP LEU LYS TRP ILE ASN ARG	GAC CTG CAG GAC TAC AAC AGA TCG CAT GGT GCA GAC TCG GAG TCG GAG ATT AAC ACA		
200	205	210	215
ASP LEU GLY ARG ARG SER ASP SER ALA MET ALA TYR VAL MET PRO THR MET ARG ASN LYS	GAC CTC GGC AGA AGA TGG CAT TCT GCG GAC GCG TAC GTC CAC CCA ACT ATC ACA AAC AAG		
220	225	230	235
CLN SER LEU PHE LEU ILE THR SER THR LYS CYS ASP LYS VAL ILE ILE GLU ASP GLY LYS	CAC AGC CTG TTC CTG ATC ACC TCG ACC AAG TGT GAC AAG CTG ATC ATC CAG GAC GGC AAG		
240	245	250	255
ALA VAL ALA VAL ARG THR VAL PRO MET LYS PRO LEU ASN PRO LYS LYS PRO VAL SER ARG	GCT GTC GCC CTG ACA ACA GTC CCA ATG AAG CCT CTG AAC GCT AAG AAC CCT GTG TCC ACA		
260	265	270	275
THR PHE ARG ALA ARG LYS GLN ILE VAL ILE SER CYS GLY THR ILE SER SER PRO LEU VAL	ATC TTC AGA GCC ACA AAG CAG ATT GTG ATC TCC TCC GCA ACC ATC TCG TCT CCT CTG GTG		
280	285	290	295
LEU GLN ARG SER GLY ILE CLY ALA ALA MET MET LEU ARG SER VAL GLY VAL LYS PRO ILE	CTC CAG ACA TCT GCT ATT GGT GCA GCT CAC CAC TTG ACA TCC GTC GGC GTC AAG CCA ATC		

**THIS PAGE BLANK (USPTO)**



300 VAL ASP LEU PRO GLY VAL GLY 305 GLU ASN PHE GLN ASP 310 HIS TYR CYS PHE PHE THR PRO THR  
 CTC GAC CTC GCA GGT CTC GGT 325 GAG AAT TTC CAC GAC CAC TAC TGT TTC TTC ACT CCA TAC  
 320 TTR VAL LYS PRO ASP VAL PRO TTR PRK ASP ASP PHE VAL ARG GLY ASP PRO VAL ALA GLR  
 TAC CTC AAG GCT GAC GTT CTT ACC TTC GAC GAC TTT CTC ACC GCG GAC CCA GTT CCC CAC  
 340 LYS ALA ALA PRE ASP GLN TRP TTR SER ASN LYS ASP GLY PRO LEU THR THR ASN GLY ILE  
 AAG GCC GCT TTC GAC CAG TCG TAC TCC AAC AAG GAC GGT CCA TTG ACC ACC AAC GGT ATT  
 360 GLU ALA GLY VAL LYS ILE ARG PRO TTR GLU GLO GLO LEU ALA THR ALA ASP GLU ASP PHE  
 GAA GCC GCA GTC AAG ATC AGA GCT ACC GAA GAC GAC CTC GCT ACC GCG GAC GAG CAC TTC  
 380 ARG ARG GLY TTR ALA GLO TTR PHE GLU ASN LYS PRO ASP LYS PRO LEU MET HIS TTR SER  
 AGA CGC GCT TAC GCA GAC TAC TTC GAG AAC AAG CCA GAC AAC CCT CTG ATG CAC TAC TCT  
 400 VAL ILE SER GLY PHE PHE GLY ASP HIS THR LYS ILE PRO ASN GLY LYS PHE MET TTR MET  
 GTC ATC TCC GCG TTC TTT GGA GAC CAC ACC AAC ATT CTT AAC GCG AAG TTC ATG ACC ATG  
 420 PHE HIS PHE LEU GLU TTR PRO PHE SER ARG CLY PRE VAL ARG ILE THR SER ALA ASN PRO  
 TTC CAC TTC CTC GAC TAT CCA TTC TCC AGA GGA TTT GTT ACA ATC ACC TCG GCA AAC CCA  
 440 TTR ASP ALA PRO ASP PHE ASP PRO CLY PHE LRU ASN ASP GLU ARG ASP LEU TRP PRO MET  
 TAC GAC GCT GCT GAC TTC GAT CCG GCG TTC CTC AAT GAC CAC CAC CAA AGA GAC CTG TCG CTT ATG  
 460 VAL TRP ALA TTR LYS LYS SER ARG GLU TTR ALA ARG ARG MET GLU SER PRE ALA GLY GLU  
 CTC TCG CCA TAC AAG AAC TCC AGA GAG AAG CCC AGA AGA ATG CAG ACC TTT GCA GGA GAG  
 480 VAL TTR SER MET HIS PRO LEU PHE LYS VAL ASP SER PRO ALA ARG ALA ARG ASP LEU ASP  
 GTC ACC TCG GAC CAC CCA TTC TTC AAC CTT GAC TCG CCA CCC AGA GCG AGA GAC CTC CAC  
 500 LEO GLU TTR CYS SER ALA TTR ALA CLY PRO LYS HIS LEU THR ALA ASN LEU TTR HIS GLY  
 CTC GAC ACA TCG ACT CCA TAT GCG GGT CTT AAC CAC CTC ACT CCC AAC CTG TAC CAC GCG  
 520 SER TRP THR VAL PRO ILE ASP LYS PRO THR PRO LYS SER ASP PHE HIS VAL THR SER ASN  
 TCG TCG ACC GTT CTT ATC CAC AAG CCA ACC CTT AAG AAC GAT TTC CAC CTG ACC TCC AAC  
 540 GLR VAL GLR LEU HIS SER ASP ILE CLU TTR THR GLU GLU ASP ASP GLU ALA ILE VAL ASN  
 CAA CTC CAA CTC CAC TCC GAC ATC CAC TAC ACC CAC GAG CAC CAC GCG ACC ATC CTC AAC  
 560 TTR ILE LYS GLU HIS THR GLU TTR TTR TRP HIS CYS LEU GLY THR CYS SER MET ALA PRO  
 TAC ATT AAG CAA CAC ACC GAC ACC ACT TCG CAC TGT CTG GGT ACC TCG TCG ATG CCC CCA  
 580 ARG GLO GLY SER LYS ILE ALA PRO LYS CLY CLY VAL LEU ASP ALA ARG LEU ASN VAL TTR  
 AGA GAC GCT ACT AAG ATT GCT GCT AAG GGA GGT CTC TTG GAC CCC AGA CTG AAC GTT TAC  
 600 CLY VAL GLR ASN LEU LYS VAL ALA ASP LEU SER VAL CYS PRO ASP ASN VAL GLY CYS ASN  
 GCA CTC CAC AAC CTC AAG GTT GCG GAC CTT TCT GTT TGT CCC GAC AAC CTT GCA TCC AAC  
 620 TTR TTR SER THR ALA LEU THR ILE GLY GLO LYS ALA ALA THR LEU VAL ALA GLU ASP LEU  
 ACC TAC TCT ACT CCA TTC ACC ATC GGT CAC AAG GCT GCC ACT CTT GTT CTT CAA GAT CTT  
 640 GLY TTR SER CLY SER ASP LEU ASP MET THR ILE PRO ASN PHE ARG LEO GLY THR TTR GLU  
 CGC TAC TCA GCG TCC CAC CTC GAC ATG ACC ATT CCA AAC TTC AGA CTC GCA ACT TAC GAG  
 660 GLU THR CLY LEU ALA ARG PHE \*\*\*  
 GAG ACC CGA CTT CCC AGA TTC TAA CCAC ACCTGCAAGC ACATACCGCT TTTCACAAGC


2000  
 GTGTTTCAAA ATAGTTCTTT TTCTGCTTTA TATCGTTTAT CAAGTGATGA CATGAAAAGC  
 2030  
 TCAAAATAGCG ACTATACGAA AATTTAATGA AAATTAATTT AAATATTTTC TTAGCGTATT  
 2100  
 ACTCACCTTC AAAATGCCCG CCGCTTCTAA CAACGTTCTC ATGATCGACA ACTACGACTC  
 2150  
 CTTTACCTGG AACCTCTACG ACTACCTCTC TCACCAGGGA CCCAATGTCG AGGTTTTTCAG  
 2200  
 GAACGATCAG ATCACCATTG CCGAGATTGA CCAGCTCAAG CCCGACCTTC TCGTCATATC  
 2300  
 CCGTGCTCTT CCGCATCCAA CAACAGACTC CGGAATATCT CCGCAGCTGA TCAGCCATT  
 2350  
 TAAACGCAAC ATTCCTGCTT TTGCTGCTCT TATGGGCCAC CACTGTATCT TCGAGGACTT  
 2400  
 TGGCGGACAC CTCGAGTATG CCGCGGAGAT TGTCCATGGA AAAACGTCCA CTGTTAAGCA  
 2450  
 CGACAACAAG CGAATCTTCA AAAACGTTCC CCAAGATGTT CCTGTACCA GATACCACTC  
 2500  
 CCTGCCCCGA ACCCTCAAGT CGCTTCCGGA CTGTCTAGAC ATCACTGCTC GCACAGACAA  
 2600  
 CCGGATCATT ATGGCTGTGA CACACAAGAA GTACACCATC GAGGCGCTCC AGTTTCATCC  
 2650  
 AGACAGCATT CTGACCGAGC ACCGCCATCT CATGATCCAG AATATCTTGA ACCTTTCCCG  
 2700  
 TGGTACTGCG CAGGAAAATG CCAACGGCGC GCCTCAGACA AAGGAAAAGCA TATTGAGAA  
 2750  
 AATATACGGG CAGAGACGAA AAGACTACGA GTTTCAGATG AACAGACCGG GCGCCAGATT  
 2800  
 TCGTATCTA GAACTGTACT TCTCCATCGG ACTGCACCGC CCCTAATCAA TTTTACGAC  
 2900  
 AGATTGCGAG ACAACATCAG CCGCGGCAAG GTTGCAATTG TCAGCGAAAT CAACAGAGCG  
 2950  
 TCGCCTTCTA AAGGCGTCTAT CGACGGAGAC GCTAACGCTG CCAACAGAGC CCTCAACTAC  
 3000  
 CCGAAGGCTG CAGTTCCCAAC AATTTCTCTT TTCACCGAGC CAACCTCGTT TAAACGAAAT  
 3050  
 ATCCAGCACC TCGAGCTCGC CACAAAAGGC ATTCACTCTG TCGGCAATAC ACCCTGTATT  
 3100  
 TTGGCGAAGC ACTTTATCTT CAACAAGTAC CAAATTCTAC ACCCCGCACT GCGCCGACCA  
 3200  
 GACACGGTTC TGCTGATTCT CAAGATGCTG AGCTC  
 3250

**THIS PAGE BLANK (USPTO)**

13/32

0173378

Fig. 12A



**THIS PAGE BLANK (USPTO)**

Fig. 10B

Promoter MOX-Neomycinphosphotransferase adaptor fragments

NEO3 5'CGGTGGTGACATCAATCTAAAGTACAAA 3'


NEO6 5'TCATTTTGTFTTTGTACTTTAGATTGATGTCACCACCGTGCA 3'

NEO7 5'AACAAAATGATTGAACAAGATGGATTGCACGCAGGTTCTCCGGCCGCTTG 3'

NEO8 5'AATTCAAGCGGCCGAGAACCTGCGTGCAATCCATCTTGTTCAA 3'

**THIS PAGE BLANK (USPTO)**

Fig. 120



**THIS PAGE BLANK (USPTO)**



16/32

PROMOTER MOX/AAO ADAPTOR----->>

-34 1

CGGTGG TGACATCAAT CTAAAGTACA AAAACAAAAT GAGAGTTGTC GTTATTGGTG  
 ACGTGCCACC ACTGTAGTTA GATTTTCATGT TTTTGTTTIA CTCTCAACAG CAATAACCAC  
 HgiAI Met

<<----->

62

CCGGTGTGAT CGGTCTGTGG ACCGCCCTGT GTATCCACGA GAGATACCAC TCCGTTCTGC  
 GGCCACAGTA GCCAGACAGC TGGCGGGACA CATAGGTGCT CTCTATGGTG AGGCAAGACG  
 SalI

122

AGCCTCTGGA CGTTAAGGTC TACGCCGACA GATTCACCCC TTTCACCACC ACCGACGTTG  
 TCGGACACCT GCAATTCACG ATGCGGCTGT CTAAGTGGGG AAAGTGGTGG TGGCTGCAAC

182

CCGCCGGTCT GTGGCAGCCT TACACCTCCG AGCCTTCCAA CCCTCAGGAG GCCAACTGGA  
 GGGGCCCAGA CACCGTCGCA ATGTGGAGGC TCGGAAGGTT GGGAGTCCTC CGGTTGACCT

242

ACCAGCAGAC CTTCAACTAC CTCCTCTCCC ACATCGGTTT GCCTAACGCC GCCAACATGG  
 TGCTCGTCTG GAAGTTGATG GAGCAGAGGG TGTAGCCAAG CGGATTGCGG CGGTTGTACC

302

GTCTGACCCC TGTCTCGGGT TACAACCTGT TCAGAGAGGC CGTTCCTGAC CTTTACTGGA  
 CAGACTGGGG ACAGAGCCCA ATGTTGGACA AGTCTCTCCG GCAAGGACTG GGAATGACCT

362

AGGACATGCT CCTCGGTTTC AGAAAGCTTA CCCCTAGAGA GCTGGACATG TTCCCTGACT  
 TCCTGTACCA GGAGCCAAAG TCTTTGGAAT GGGGATCTCT CGACCTGTAC AAGGGACTGA  
 HindIII

422

ACAGATACGG TTGGTTCAAC ACCTCCCTGA TCCTGGAGGG TAGAAAGTAC CTGCAGTGGC  
 TGCTATATGCC AACCAGTTG TGGAGGGACT AGGACCTCCC ATCTTTCATG GACGTCACCG

482

TGACCCGAGC ACTGACCGAG AGAGGTGTTA AGTTCTTCTT GAGAAAGGTC GAGTCCTTCG  
 ACTGGCTCTC TGACTGGCTC TCTCCACAAT TCAAGAAGGA CTCTTTCACG CTCAGGAAGC

542

AGGAGGTTGC CAGAGGTGGT GCGGACGTCA TCATCATGTG TACCGGTGTC TGGGCCGGTG  
 TCCTCCAACG GTCTCCACCA CGGCTGCACT AGTAGTACAC ATGGCCACAG ACCCGGCCAC

602

TCCTGCAGCC TGACCTCTCT CTGCAGCCCG GGAGAGGTCA GATCATTAAAG GTTGACGCCC  
 AGGACGTCGG ACTGGGAGAC GACGTCGGGC CCTCTCCAGT CTAGTAATTC CAACTGCGGG  
 XbaI

662

CATGGCTGAA GAACCTCATC ATTACCCACG ACCTGGAGAG AGGTATCTAC AACTCCCTT  
 GTACCGACTT CTGAAGTAG TAATGGGTGC TGGACCTCTC TCCATAGATG TTGAGGGGAA

722

ACATTATCCC TGGTCTGCAG GCCGTACCCC TGGGTGGTAC CTTCCAGGTC GGTAACCTGA  
 TGTAAATAGG ACCAGACGTC CGGCAGTGGG ACCCACCATG GAAGGTCCAG CCATTGACCT  
 KpnI

782

ACGAGATCAA CAACATCCAG GACCACAACA CCATCTGGGA GGGTTGTTGT AGACTGGAGC  
 TGCTCTAGTT GTGTAGGTC CTGGTGTGTT GGTAGACCCT CCCAACAACA TCTGACCTCG

842

CTACCCTGAA GGACGCCAAG ATCGTTGGTG AGTACACCGG TTTCAGACCT GTTAGACCTC  
 GATGGGACTT CCTGCGGTTT TAGCAACCAC TCATGTGGCC AAAGTCTGGA CAATCTGGAG

902

AGGTCAGACT GGAGAGAGAG CAGCTGAGAT TCGGTTCTCT CAACACCGAG GTCATTACCA  
 TCCAGTCTGA CCTCTCTCTC GTCGACTCTA AGCCAAGGAG GTTGTGGCTC CAGTAAGTGT

962

ACTACGGTCA CGGTGGTTAC GGTCTGACCA TCCACTTGGG TTGTGCCCTG GAGGTTGCCA  
 TGATGCCAGT GCCACCAATG CCAGACTGGT AGGTGAACCC AACACGGGAC CTCCAACGGT


1022

AGCTGTTCCG TAAGGTCCTG GAGGAGAGAA ACCTGCTGAC CATGCCCTCA TCCCACCTGT  
 TCGACAAGCC ATTCCAGGAC CTCCTCTCTT TGGACGACTG GTACGGAGCT AGGGTGGACA

GAG  
 CTCAGCT  
 \*\*SalI


**THIS PAGE BLANK (USPTO)**

FIG. 14A



**THIS PAGE BLANK (USPTO)**

Fig. 14B




**THIS PAGE BLANK (USPTO)**

19/32

0173378

Fig. 14C



**THIS PAGE BLANK (USPTO)**



20/32

0173378

<-----PROMOTER MOX-HGRF ADAPTOR----->>  
-34

CGGTG GTGACATCAA TCTAAAGTA CAAAAACAAA  
ACGTGCCAC CACTGTAGTT AGATTTCAT GTTTTGTITT  
HgiAI

<<----->>  
1

ATGTACGCCG ACGCCATCTT CACCAACTCC TACAGAAAGG TTCTGGGTCA GCTCTCGGCC  
TACATGCGGC TCGGGTAGAA GTGGTTGAGG ATGTCTTTCC AAGACCCAGT CGAGAGCCGG  
Met

--->  
61

AGAAAGCTTC TGCAGGACAT CATGTCGAGA CAGCAGGGTG AGTCCAACCA GGAGAGAGGT  
TCTTTCGAAG ACGTCCTGTA GTACAGCTCT GTCGTCCCAC TCAGGTGGT CCTCTCTCCA  
HindIII PstI

121

GCCAGAGCCA GACTGTGAG  
CGGTCTCGGT CTGACACTCA GCT  
\*\*\* Sali


Fig. 15

**THIS PAGE BLANK (USPTO)**

21/82


0173378

Fig. 16A




**THIS PAGE BLANK (USPTO)**

Fig. 16B



**THIS PAGE BLANK (USPTO)**


Fig. 16C



**THIS PAGE BLANK (USPTO)**




Fig. 16D




**THIS PAGE BLANK (USPTO)**

Fig. 16E



**THIS PAGE BLANK (USPTO)**

Fig. 16F



**THIS PAGE BLANK (USPTO)**

1 CATGTACGCCG ACGCCATCTT CACCAACTCC TACAGAAAGG TTCTGGGTCA GCTCTCGGCC  
 CATGGTACATGCCGC TCGGGTAGAA GTGGTTGAGG ATGCTTTTCC AAGACCCAGT CGAGAGCCCGG  
KpnI Met

61 AGAAAGCTTC TGCAGGACAT CTGTTGAGA CAGCAGGGTG AGTCCAACCA GGAGAGAGGT  
 TCTTTCGAAG ACGTCTGTA GACAAGCTCT GTCGTCCCAC TCAGGTGGT CCTCTCTCCA  
HindIII PstI cys

121 GCCAGAGCCA GACTGTGAGGTAC  
 CGGTCTCGGT CTGACACTC  
 \*\*\* KpnI

Fig. 17

**THIS PAGE BLANK (USPTO)**



Fig. 18A

CTTCGCCAAT GATTCAGCTG CTGGACCGAA AACGCCCTCTT TTGGCCAAAAA AAAGCCCACC  
 -2104  
 GTTGATAACT GCGGAGGCCA TATTTCAAAG AACAGCGAAT AAAAAAAAAA GGTGAATGAA  
 -2054 -2004  
 ATGCGCGAAA CGATACCACT TATTAGCATA AACAAAAAAA AAAAAAATCT ATTAGCTGTT  
 -1954  
 ATTATAATTA GTTCAATAAT TTCATAAGCA TCATCGTTGG GCGGCCTATT GTCATCAGTG  
 -1904  
 GTCCCTCTGG AACAGGTAAA TCCACTTTGC TGAAGAAGCT GTTTGCTGAG TTCCAGACA  
 -1854  
 AGTTTGGATT TTCCGTGTCC AACACCACGA GAAAACCTAG ACCTGGTGAA AAAGACGGTG  
 -1804  
 TCGATTACCA CTTACCACAG GTAGAGGACT TCAAGAAGAT GATTGAAGAA AACAAATTCA  
 -1754 -1704  
 TTGAATGGGC CCAGTTCTCC GGCAACTACT ACGGCACCTC TGTGAAAGCT GTGCAAGACG  
 -1654  
 TGGCCGAAGT GATGAAGAGA ACGTGTATTT TGCACATTGA TATGCAGGGT GTCAAGAGCG  
 -1604  
 TCAAGAAGAC CAACCTGGCA CCCCCATTCC TCTTTATTTT TCCTCCGTCC ATCGAAGAGC  
 -1554  
 TCAAGAAGAG GCTCGAGAGC CGTGGAACAG AGACCCCTGA ATCTCTTGCC AAGCGGCTTG  
 -1504  
 CTGCTGCATC TGCGGAGATC GAGTACGCCA GGGCAGTGGA CACGACAAGG TCATTGTCAA  
 -1454 -1404  
 CGATGACCTT GAGAAGCGCT ACTCTGAGCT GAAGGAGTTC ATTTTCGCCG AGCCCATCTA  
 -1354  
 AGCATTGATA AATTTTTAAT ATCTAGAGCT CTCATACGGG ACAGTATCTC CTCCAACCTT  
 -1304  
 CCGTCAAGCT TGTCCTCTTC ATGCTCCTCA ACAGTCATGG CATCCAGCTG CTGCTGCTTT  
 -1254  
 TGCTCCAGCC TGGCATATAT GTCGCCATAC AGCTTGAGTT GGATTTTGAT GAAACTCTCA  
 -1204  
 AAGGTAGGGT CCACCAGTGA CAGTCGCAGC GCAATGAACT GCTCGATTTC GTTCTTGAGC  
 -1154 -1104  
 CGTGTGTTGA TGTCCGTGTA GATATTTTCT GCCTCGTCGT ACTCAACTTT GAACTTCTGC  
 -1054  
 AGCTTGTTCA GGCTCTTCTG TAACTGGTCT GTTTTCTCGG TGTGATGCTG CTCGGTCACC  
 -1004  
 TGTCGGTCAA TCGCTTCGTA CTCGGCTCTG AGCTTCGAAA GCTTGAATCG TGAACGTCG  
 -954  
 TAATCCACCT TTTTCCGTGC GCGCTTCTTG ATCAGCTTGT TGATCTCGTC GTTGTACTTC  
 -904  
 TTCAGCTCGT TAATCGGCTC CACGACCGTC ATGCTCATTG GCTCCAGAAT TTCTGGCAGA  
 -854 -804  
 ATATTGTCTT TGATGTCTTC CACCATCTGC AGATAATTCA GAGAAATACC ATCTCTGGGG  
 -754  
 TTCACCTTGT GCTCTTCTGG CCGTTCCGCA GCTTCCGACC GCTTATCAGC CTTGAGCTCA  
 -704  
 AAGCTATAGT CTCCGTAAAA CGACTCCAGT GTTCTAGCCA TATTTATCTG AGTCTCGAGC  
 -654  
 AGATTCTCCG AAATTGCCCA CAAAACGGCC TAGTTCCTGG TCCAGCTCGT TGGTGTAAGT  
 -604  
 CTCGAGTTTG CGGAAATTGG CCTCCTGGAC GTCAAACTCA GGATCAACAG AGGGCTCACC  
 -554 -504  
 TTTGTTTGTG CGTAGTATCA CATGTGCTCC GGCACGATTG ACAGCTTTTT TAAACCCAAC  
 -454  
 CCATGACATG TCGAGGAAAG GGTCTTTTCG GCGAGTTAAA TATTTTGGC TATGTAGCAG  
 -404  
 ACATGTTTCG ACGCTGCCGT CCGCTCGATC GGAAAATATT ACCCCAGGAA CAAGCACTTG  
 -354  
 CTTGGGTTAG CCACCACCTT GCGCAAGCCT TTTTGCCGGC TCTACACAGG GCCAATGAAA  
 -304  
 TCTGGCGCGA ATCTGAAACC CATGAAACGG ACGACACTGC CAACAAGCTC ACTGCACTAT  
 -254 -204

**THIS PAGE BLANK (USPTO)**

29/32

0173378

Fig. 183

TTTTTCCTC TAGTGAATA GCCTATCCTC GTCTCGCTCC CCTCATACCT GTAAAGGGGT  
 -154  
 GCAATTTAGC CTCGTTCCAG CCATTACGGG GCCACTCAAC AACACGTCCG CTACCATGGG  
 -104  
 GTGCTTGGGC ACCAAAAGGC CTATAAATAG GCCCCCATCC GTCTGCTACA CAGTCATCTC  
 -54

1	5	10	15
TGTCCTTTCTTCCC	MET SER MET ARG ILE PRO LYS ALA ALA	SER VAL ASN ASP GLU GLN HIS	
-14	ATG AGT ATG AGA ATC CCT AAA GCA GCG	TCG GTC AAC GAC GAA CAA CAC	
20	25	30	35
GLN ARG ILE ILE LYS TYR GLY ARG	ALA LEU VAL LEU ASP ILE VAL GLU GLN TYR GLY GLY		
CAG ACA ATC ATC AAG TAC GGT CGT	GCT CTT GTC CTG GAC ATT GTC GAG CAG TAC GCA GGA		
40	45	50	55
GLY HIS PRO GLY SER ALA MET GLY	ALA MET ALA ILE GLY ILE ALA LEU TRP LYS TYR THR		
GGC CAC CCG GGC TCG GCC ATG GGC	GCC ATG GCT ATC GGA ATT GCT CTG TGG AAA TAC ACC		
60	65	70	75
LEU LYS TYR ALA PRO ASN ASP PRO	ASN TYR PHE ASN ARG ASP ARG PHE VAL LEU SER ASN		
CTG AAA TAT GCT CCC AAC GAC CCT	AAC TAC TTC AAC AGA GAC AGG TTT GTC CTG TCG AAC		
80	85	90	95
GLY HIS VAL CYS LEU PHE GLN TYR	ILE PHE GLN HIS LEU TYR GLY LEU LYS SER MET THR		
GCT CAC CTG TGT CTG TTC CAG TAT	ATC TTC CAG CAC CTG TAC GGT CTC AAC TCG ATG ACC		
100	105	110	115
MET ALA GLN LEU LYS SER TYR HIS	SER ASN ASP PHE HIS SER LEU CYS PRO GLY HIS PRO		
ATG GCG CAG CTG AAG TCC TAC CAC	TCG AAT GAC TTC CAC TCG CTG TGT CCC GGT CAC CCA		
120	125	130	135
GLU ILE GLU HIS ASP ALA VAL GLU	VAL THR THR GLY PRO LEU GLY GLN GLY ILE SER ASN		
GAA ATC GAG CAC GAC GCC GTC GAG	GTC ACA ACG GGC CCG CTC GCC CAG GGT ATC TCG AAC		
140	145	150	155
SER VAL GLY LEU ALA ILE ALA THR	LYS ASN LEU ALA ALA THR TYR ASN LYS PRO GLY PHE		
TCT GTT GGT CTG GCC ATA GCC ACC	AAA AAC CTG GCT GCC ACG TAC AAC AAG CCG GGC TTT		
160	165	170	175
ASP ILE ILE THR ASN LYS VAL TYR	CYS MET VAL GLY ASP ALA CYS LEU GLN GLU GLY PRO		
GAT ATC ATC ACC AAC AAG GTG TAC	TGC ATG GTT GCC GAT GCG TGC TTC CAG GAG GCC CCT		
180	185	190	195
ALA LEU GLU SER ILE SER LEU ALA	GLY HIS MET GLY LEU ASP ASN LEU ILE VAL LEU TYR		
GCT CTC GAG TCG ATC TCG CTG GCC	GCC CAC ATG GGG CTG GAC AAT CTG ATT GTG CTC TAC		
200	205	210	215
ASP ASN ASN GLN VAL CYS CYS ASP	GLY SER VAL ASP ILE ALA ASN THR GLU ASP ILE SER		
GAC AAC AAC CAG GTC TGC TGT GAC	GCC AGT GTT GAC ATT GCC AAC ACC GAG GAC ATC AGT		
220	225	230	235
ALA LYS PHE LYS ALA CYS ASN TRP	ASN VAL ILE GLU VAL GLU ASN ALA SER GLU ASP VAL		
GCC AAG TTC AAG GCC TGC AAC TGG	AAC GTG ATC GAG GTC GAG AAC GCT TCC CAG GAC GTC		
240	245	250	255
ALA THR ILE VAL LYS ALA LEU GLU	TYR ALA GLN ALA GLU LYS HIS ARG PRO THR LEU ILE		
GCT ACC ATT GTC AAG GCC TTC CAG	TAC GCG CAG GCC GAG AAC CAC AGA CCA ACA CTT ATC		
260	265	270	275
ASN CYS ARG THR VAL ILE GLY SER	GLY ALA ALA PHE GLU ASN HIS CYS ALA ALA HIS GLY		
AAC TGC AGA ACT CTG ATT GGA TCG	GCT GCT GCG TTC GAG AAC CAC TGT GCT GCG CAC GGT		
280	285	290	295
ASN ALA LEU GLY GLU ASP GLY VAL	ARG GLU LEU LYS ILE LYS TYR GLY MET ASN PRO ALA		
AAC GCT CTG GCC CAG CAC GCT CTG	CGC GAG CTC AAA ATC AAG TAC GCG ATG AAC CCG GCC		
300	305	310	315
GLN LYS PHE TYR ILE PRO GLN ASP	VAL TYR ASP PHE PHE LYS GLU LYS PRO ALA GLU GLY		
CAG AAG TTC TAC ATT CCG CAG GAC	GTC TAC GAC TTC TTC AAG CAG AAG CCG GCC GAG GGC		
320	325	330	335
ASP LYS LEU VAL ALA GLU TRP LYS	SER LEU VAL ALA LYS TYR VAL LYS ALA TYR PRO GLU		
GAC AAG CTG CTG GCC GAA TGC AAG	AGT CTC GTG GCC AAG TAC GTC AAC CCG TAC CCT CAG		
340	345	350	355
GLU GLY GLN GLU PHE LEU ALA ARG	MET ARG GLY GLU LEU PRO LYS ASN TRP LYS SER PHE		
GAG GGC CAG GAG TTT TTC GCG CCG	ATG ACA GGC GAG CTG CCA AAG AAC TGG AAG TCG TTC		
360	365	370	375

**THIS PAGE BLANK (USPTO)**

0173378

Fig. 18C

LEU	PRO	GLN	GLU	GLU	PHE	THR	GLY	ASP	ALA	PRO	THR	ARG	ALA	ALA	ALA	ARG	GLU	LEU	VAL
CTG	CCG	CAG	CAG	GAA	TTC	ACC	GGC	GAC	GCT	CCT	ACA	AGG	GCC	GCT	GCC	AGA	GAG	CTT	GTG
			380					385					390					395	
ARG	ALA	LEU	GLY	GLN	ASN	CYS	LYS	SER	VAL	ILE	ALA	GLY	CYS	ALA	ASP	LEU	SER	VAL	SER
AGA	GCC	CTG	GGG	CAG	AAC	TGC	AAG	TCG	GTG	ATT	GCC	GGT	TGC	GCA	GAC	CTG	TCT	GTG	TCT
			400					405					410					415	
VAL	ASN	LEU	GLN	TRP	PRO	GLY	VAL	LYS	TYR	PHE	MET	ASP	PRO	SER	LEU	SER	THR	GLN	CYS
GTC	AAT	TTG	CAG	TGG	CCA	GGG	GTG	AAA	TAT	TTC	ATG	GAC	CCC	TCG	CTG	TCC	ACG	CAG	TGT
			420					425					430					435	
GLY	LEU	SER	GLY	ASP	TYR	SER	GLY	ARG	TYR	ILE	GLU	TYR	GLY	ILE	ARG	GLU	HIS	ALA	MET
GGC	CTG	AGC	GGC	GAC	TAC	TCC	GGC	AGA	TAC	ATT	GAG	TAC	GGA	ATC	AGA	GAA	CAC	GCC	ATG
			440					445					450					455	
CYS	ALA	ILE	ALA	ASN	GLY	LEU	ALA	ALA	TYR	ASN	LYS	GLY	THR	PHE	LEU	PRO	ILE	THR	SER
TGT	GCT	ATC	GCC	AAT	GGC	CTT	GCC	GCC	TAC	AAC	AAG	GGC	ACG	TTC	CTG	CCG	ATC	ACG	TCC
			460					465					470					475	
THR	PHE	PHE	MET	PHE	TYR	LEU	TYR	ALA	ALA	PRO	ALA	ILE	ARG	MET	ALA	GLY	LEU	GLN	GLU
ACT	TTC	TTC	ATG	TTC	TAC	CTG	TAC	GCT	GCC	CCA	GCC	ATC	AGA	ATG	GCC	GGC	CTG	CAG	GAG
			480					485					490					495	
LEU	LYS	ALA	ILE	HIS	ILE	GLY	THR	HIS	ASP	SER	ILE	ASN	GLU	GLY	GLU	ASN	GLY	PRO	THR
CTC	AAG	GCG	ATC	CAC	ATC	GGC	ACC	CAC	GAC	TCG	ATC	AAT	GAG	GGT	GAG	AAC	GGC	CCT	ACG
			500					505					510					515	
HIS	GLN	PRO	VAL	GLU	SER	PRO	ALA	LEU	PHE	ARG	ALA	TYR	ALA	ASN	ILE	TYR	TYR	MET	ARG
CAC	CAG	CCG	GTC	GAG	TCG	CCA	GCA	TTG	TTC	CGG	GCC	TAT	GCA	AAC	ATT	TAC	TAC	ATG	AGA
			520					525					530					535	
PRO	VAL	ASP	SER	ALA	GLU	VAL	PHE	GLY	LEU	PHE	GLN	LYS	ALA	VAL	GLU	LEU	PRO	PHE	SER
CCG	GTC	GAC	TCT	GCA	GAA	GTG	TTT	GGC	CTG	TTC	CAA	AAA	GCC	GTC	GAG	CTG	CCA	TTC	AGC
			540					545					550					555	
SER	ILE	LEU	SER	LEU	SER	ARG	ASN	GLU	VAL	LEU	GLN	TYR	LEU	ALA	SER	ARG	ALA	GLN	ARG
TCG	ATT	CTG	TCG	CTC	TCG	AGA	AAC	GAG	GTG	CTG	CAA	TAC	CTG	GCA	AGT	CGA	GCG	CAG	AGA
			560					565					570					575	
ARG	ARG	ASN	ALA	ALA	GLY	TYR	ILE	LEU	GLU	ASP	ALA	GLU	ASN	ALA	GLU	VAL	GLN	ILE	ILE
AGG	CGC	AAC	GCG	GCC	GGC	TAT	ATT	CTG	GAG	GAT	GCG	GAG	AAC	GCC	GAG	GTG	CAG	ATT	ATT
			580					585					590					595	
GLY	VAL	GLY	ALA	GLU	MET	GLU	PHE	ALA	ASP	LYS									

**THIS PAGE BLANK (USPTO)**



**THIS PAGE BLANK (USPTO)**



Fig. 20Identical sequences in -1000 region of DAS and MOX genes

DAS -1076

TAGATATTTTCTGCCTCGTCGTA

\* \*\* \*\*\*\*\* \* \*\*\*\*\* \*

\* \*\* \*\*\*\*\* \*

TCGAAATTTTGGCGTCGTACAGTGTGATGTCACC

MOX -1052

DAS

-937

ATCGCTTCGTA

\*\*\*\*\* \* \* \* \*

\*\*\*\*\* \* \* \*

ATCGAATGTAATGAGCTGCAGCTTGCGA

MOX

-987

**THIS PAGE BLANK (USPTO)**



European Patent Office

Application number: 0173378

85201235.0

**DECLARATION PURSUANT TO RULE 28, PARAGRAPH 4,  
OF THE EUROPEAN PATENT CONVENTION**

The applicant has informed the European Patent Office that, until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, the availability of the micro-organism(s) identified below, referred to in paragraph 3 of Rule 28 of the European Patent Convention, shall be effected only by the issue of a sample to an expert.

**IDENTIFICATION OF THE MICRO-ORGANISMS**

Accession numbers of the deposits:

CBS 7171

CBS 7172

ATCC 34438

**THIS PAGE BLANK (USPTO)**



0173378

# Unilever N.V.

Patent Division  
PO Box 137  
3130 AC Vlaardingen  
The Netherlands

Olivier van Noortlaan 120  
Tel. 010-60 69 33, (UTN Code 322)  
Telex 23261, Fax (gr II) 605800  
Cables: Unileverpatent



European Patent Office  
P.O. Box 5818  
2280 HV Rijswijk

PLEASE ENDORSE AND  
RETURN DUPLICATE COPY  
AS ACKNOWLEDGEMENT.



Your Ref

Our Ref JVT/PS

Direct Dial. 60 . . . 5741

Date

29 Aug. 1985

Re.: Recently filed patent application No. 85201235.0 - our case  
T 7000 (R)-EP

In this patent specification a Fig. 3 was present giving the complete nucleotide sequence of the HARS-1 fragment (see page 58). This sequence was determined shortly before the expiry of the priority year. Re-analysis of the experimental data has revealed that the sequence contained several errors.

A corrected sequence of the complete nucleotide sequence of the HARS-1 fragment is now provided.

It is requested that this correction of errors made by Applicants is allowed by the Patent Office in order to correct a part of the disclosure which is now known to be wrong.

Van der Toorren, Johannes Drs.  
European Patent Attorney  
General Authorization No. 170

**THIS PAGE BLANK (USPTO)**

Fig. 3 (amended)

DNA sequence of the autonomous-replicating sequence HARS1 from the methylotrophic yeast Hansenula polymorpha. The HARS1 represents a SalI fragment comprising 499 nucleotides. The dideoxy-sequencing method was employed.

<sup>2</sup>  
 ↑  
 (G)TCGACTCCC GCGACTCGGC GTTCACTTTC GAGCTATTAT 40  
 CAACGCCGGA ATACGTCAGA AACAGCCGTG CCCCAGGGAC 80  
 CAGAAAGCCT ACTGGTGAGT ATGTTCTTTC GTGTGATTTT 120  
 TCCGAGGATG AGAACGACGA TAACGAGCAC AACTCGGAGT 160  
 CGGAGGACAC GCTTATTGCG TTGAACGCAG CCACATCAGC 200  
 AGGCTGTCAA GACTGAGTAT GGCCACAGAG CTGGATTCTC 240  
 GGCCTCATAC TCAAGACGTT AGTAAACTCC GTCTGCCAGA 280  
 AATTGCTGAC GAGGATGTAT AATAATAGAT GAATTACGAA 320  
 CAATTGTAGT TCAAAAAAAT TTAGTAACAA TATTGTCTAG 360  
 ATGACAGATG TGCTGAAACC AGTGA ACTCC AATAAACCAC 400  
 TCACCGCTAC CCAAGAGAAA CAGATCAGAG TGCTAGGGCC 440  
 TTGTTTCAGA GTACTACAAC GTTTACCAGA AGCTTGAGCA 480  
 AGTTCTCAA CGCGGGTTTG (TCGAC)

↓  
500

**THIS PAGE BLANK (USPTO)**



EP 0173378 (2)

~~C12N15/26~~ (C12N15/78) - (C12N15/63) -  
(C12N15/82) - (C12N15/22) - (C12N15/66) -  
(C12N9/00) - (C12N9/04) - (C12N9/06) -  
(C11D3/386) -

-10-BASIC DOC.-



Office européen des brevets

(11) Publication number:

0 173 378

A3

C12N15/53

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 85201235.0

(22) Date of filing: 25.07.85

(51) Int. Cl.<sup>4</sup>: C 12 N 15/00

C 12 N 1/16, C 11 D 3/386  
C 11 D 3/395, C 12 P 21/02  
C 07 H 21/04, C 12 N 9/02

A request for correction of Fig. 3 has been filed pursuant to Rule 88 EPC. A decision on the request will be taken during the proceedings before the Examining Division.

(30) Priority: 27.07.84 EP 84201114  
07.02.85 GB 8503160

(43) Date of publication of application:  
05.03.86 Bulletin 86/10

(88) Date of deferred publication of search report: 06.08.86

(84) Designated Contracting States:  
AT BE CH DE FR GB IT LI NL SE

(71) Applicant: UNILEVER NV  
Burgemeester s'Jacobplein 1 P.O. Box 760  
NL-3000 DK Rotterdam(NL)

(84) Designated Contracting States:  
BE CH DE FR IT LI NL SE AT

(71) Applicant: UNILEVER PLC  
Unilever House Blackfriars P.O. Box 68  
London EC4P 4BQ(GB)

(84) Designated Contracting States:  
GB

(72) Inventor: Ledebor, Adrianus Merinus  
Burg. leFèvre de Montignyplein 8  
NL-3055 NL Rotterdam(NL)

(72) Inventor: Maat, Jan  
Molenstraat 2  
NL-2681 BS Monster(NL)

(72) Inventor: Verrips, Cornelis Theodorus  
Hagedoorn 18  
NL-3142 KB Maassluis(NL)

(72) Inventor: Visser, Christiaan  
Minoeserf 77  
NL-2907 VJ Capelle a/d IJssel(NL)

(72) Inventor: Janowicz, Zbigniew Alojzy  
Adalbert-Stifter-Strasse 49  
D-4006 Erkrath-Unterfeldhaus(DE)

(72) Inventor: Hollenberg, Cornelis Petrus  
Chopin-Strasse 7  
D-4000 Düsseldorf(DE)

(74) Representative: van der Toorren, Johannes, Drs. et al,  
UNILEVER N.V. Patent Division Postbus 137  
NL-3130 AC Vlaarding(en)(NL)

(54) Use of oxidoreductases in bleaching and/or detergent compositions and their preparation by microorganisms engineered by recombinant DNA technology.

(57) The structural genes and their regulatory DNA sequences of an alcohol oxidase (MOX) and a dihydroxyacetone synthase (DHAS) of *Hansenula polymorpha* have been isolated and the nucleotide sequences determined. The invention relates to the use of the MOX gene, as well as the use of the regulatory DNA sequences of MOX and/or DAS in combination with the MOX gene, optionally after modification thereof, or other oxidase genes, or other genes, to produce engineered microorganisms, in particular yeasts.

Said engineered microorganisms can produce oxidases or other enzymes in yields that allow industrial application on a large scale.

Moreover, said engineered microorganisms can produce oxidases having improved properties with respect to their application in oxidation reactions and/or in bleaching and detergent products.

EP 0 173 378 A3



European Patent  
Office

# EUROPEAN SEARCH REPORT

0173378

Application number

EP 85 20 1235

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	EP-A- 0 103 887 (AMGEN) * Claims 1-28 * ---	1	C 12 N 15/00 C 12 N 1/16 C 11 D 3/395 C 11 D 3/386
X	EP-A- 0 098 533 (NABISCO BRANDS) * Claims 1-7; page 4 * ---	1	C 12 P 21/02 C 07 H 21/04 C 12 N 9/02
X	EP-A- 0 086 139 (TRANSGENE) * Claims 1-38 * ---	1	
X	EP-A- 0 066 994 (ICI) * Claims 1-17 * ---	1	
X	CHEMICAL ABSTRACTS, vol.100, no.7, February 1984, page 147, ref.nr. 46308t; Columbus, Ohio, US G. BRANLANT et al.: "Molecular cloning of the glyceraldehyde-3-phosphate dehydrogenase genes of Bacillus stearothermophilus and Escherichia coli, and their expression in Escherichia coli." & GENE 1983, 25(1), 1-7. * Abstract * ---	1	TECHNICAL FIELDS SEARCHED (Int. Cl. 4)  C 12 N C 11 D
X	CHEMICAL ABSTRACTS, vol.96, no.5, March 1982, page 121, ref.no. 63538j; Columbus, Ohio, US C.A. LEE et al.: "Plasmid-directed synthesis of enzymes required for D-mannitol transport and utilization in Escherichia coli." ./.		
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of the search 17-04-1986	Examiner DELANGHE
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			



European Patent  
Office

# EUROPEAN SEARCH REPORT

0173378

Application number

EP 85 20 1235

- 2 -

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	& PROC. NATL. ACAD. SCI. USA 1981, 78(12), 7336-40.  * Abstract *  ---	1	
X	CHEMICAL ABSTRACTS, vol.94, no.21, May 1981, page 375, ref.no. 170905g; Columbus, Ohio, US E. LOHMEIER et al.: "Cloning and expression of the fumarate reductase gene of Escherichia coli." & CAN. J. BIOCHEM. 1981, 59(3), 158-64.  * Abstract *  ---	1	
X	JOURNAL OF BIOCHEMISTRY, vol.91, no.4, April 1982, pages 1205-1212; Tokyo, JP M. IWAKURA et al.: "Cloning of dihydrofolate reductase gene of Escherichia coli K12."  * Pages 1205-1211 *  ---	1	
A	CHEMICAL ABSTRACTS, vol.99, no.1, July 4, 1983, page 227, ref.no. 2206q; Columbus, Ohio, US M.J. WAITES et al.: "Dihydroxyacetone synthase: a special transketolase for formaldehyde fixation from the methylotrophic yeast Candida boidinii CBS 5777." & MICROBIOL. 1983, 129(4), 935-44.  * Abstract *  ---	1, 35	
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
<b>CATEGORY OF CITED DOCUMENTS</b>			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons  & : member of the same patent family, corresponding document	



## CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

## X LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions,

namely: 1) Claims 1-30: Process for preparing oxidoreductases by recombinant technology, their use, DNA sequences encoding oxidoreductases, process for preparing a transformed microorganism, microorganisms used

2) Claims 35-39: DNA sequences coding for DHAS

3) Claims 31-34, 40-47: DNA sequences containing a regulon and a structural gene coding for a specific enzyme or other protein. Process for preparing this enzyme.

- ☒ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☐ None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
P, X	gene coding for methanol oxidase in Hansenula polymorpha."  * Whole document. *  ---	1-30	
	NUCLEIC ACIDS RESEARCH, vol.13, no. 9, May 1985, pages 3043-3062; Oxford, GB Z.A. JANOWICZ et al.: "Cloning and characterization of the DAS gene encoding the major methanol assimilatory enzyme from the methylotrophic yeast Hansenula polymorpha."  * Whole document *  -----	1-39	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
Place of search		Date of completion of the search	Examiner
<b>CATEGORY OF CITED DOCUMENTS</b>			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons  & : member of the same patent family, corresponding document	



European Patent  
Office

# EUROPEAN SEARCH REPORT

0173378

Application number

EP 85 20 1235

- 3 -

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	CHEMICAL ABSTRACTS, vol.96, no.13, March 29, 1982, page 560, ref.no. 102478m; Columbus, Ohio, US M. BRAVO et al.: "Enzymic oxidation of methanol to produce formaldehyde and hydrogen peroxide." & ADV. BIOTECHNOL., (PROC. INT. FERMENT. SYMP.), 6th 1980 (Pub. 1981), 3, 329-34.  * Abstract *	1	
A	CHEMICAL ABSTRACTS, vol.95, no.5, August 3, 1981, page 291, ref.no. 37855v; Columbus, Ohio, US J. GEISSLER et al.: "Yeast methanol oxidases: An unusual type of flavo-protein." & FEBS LETT. 1981, 126(2), 152-6.  * Abstract *	1	
P,X	WO-A- 84 04 539 (TECHNISCHE HOGE-SCHOOL DELFT)  * Claims *	1	
P,X	WO-A- 85 01 063 (TECHNISCHE HOGE-SCHOOL DELFT)  * Claims *	1	
P,X	NUCLEIC ACIDS RESEARCH, vol.13, no. 9, May 1985, pages 3063-3082; Oxford, GB A.M. LEDEBOER et al.: "Molecular cloning and characterization of a		
The present search report has been drawn up for all claims ./. . .			
Place of search		Date of completion of the search	Examiner
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**